



Exponential stability of a network of serially connected Euler–Bernoulli beams

Denis Mercier, Virginie Régnier

► To cite this version:

Denis Mercier, Virginie Régnier. Exponential stability of a network of serially connected Euler–Bernoulli beams. *International Journal of Control*, Taylor & Francis, 2014, 87 (6), pp.1266-1281. 10.1080/00207179.2013.874597 . hal-03163679

HAL Id: hal-03163679

<https://hal-uphf.archives-ouvertes.fr/hal-03163679>

Submitted on 28 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Exponential stability of a network of serially connected Euler-Bernoulli beams

D. Mercier, V. Régnier *

December 9, 2013

Abstract

The aim is to prove the exponential stability of a system modelling the vibrations of a network of N Euler-Bernoulli beams serially connected. Using a result due to K. Ammari and M. Tucsnak, the problem is reduced to the estimate of a transfer function and the obtention of an observability inequality. The solution is then expressed in terms of Fourier series so that one of the sufficient conditions for both the estimate of the transfer function and the observability inequality is that the distance between two consecutive large eigenvalues of the spatial operator involved in this evolution problem is superior to a minimal fixed value. This property called spectral gap holds. It is proved using the exterior matrix method due to W. H. Paulsen. Two more asymptotic estimates involving the eigenfunctions are required. They are established using an adequate basis.

Key words Network, Beams, Stability, Spectral gap, Exterior matrices.

AMS 34B45, 74K10, 93B60, 93D15.

1 Introduction

In the last few years various physical models of multi-link flexible structures consisting of finitely many interconnected flexible elements such as strings, beams, plates, shells have been mathematically studied. See the references by Ali Mehmeti, von Below and Nicaise in [16] as well as [10] and [13], for instance. The spectral analysis of such structures has some applications to control or stabilization problems (cf. [13]).

For interconnected strings (corresponding to a second-order operator on each string), a lot of results have been obtained: the asymptotic behaviour of the eigenvalues (see the references by Ali Mehmeti, von Below and Nicaise in [16]), the relationship between the eigenvalues and

*Laboratoire de Mathématiques et ses Applications de Valenciennes, FR CNRS 2956, Institut des Sciences et Techniques de Valenciennes, Université de Valenciennes et du Hainaut-Cambrésis, Le Mont Houy, 59313 VALENCIENNES Cedex 9, FRANCE, e-mails : denis.mercier@univ-valenciennes.fr ; virginie.regnier@univ-valenciennes.fr

algebraic theory (cf. papers by von Below, Nicaise and [13]), qualitative properties of solutions (see papers by von Below cited in [16] for example) etc...

For interconnected beams (corresponding to a fourth-order operator on each beam), some results on the asymptotic behaviour of the eigenvalues and on the relationship between the eigenvalues and algebraic theory were obtained by Nicaise and Dekoninck with different kinds of connections using the method developed by von Below in [8] to get the characteristic equation associated to the eigenvalues.

The authors used the same method in [15] to compute the spectrum for a hybrid system of N flexible beams connected by n vibrating point masses. This type of structure was studied by Castro and Zuazua in many papers (see [9] and the papers by the same authors cited in [16] as well one by Castro and Hansen also cited there).

In another paper (see [16]), the authors used the technique of exterior matrices due to W. H. Paulsen (presented for other purposes in [18]) which D. Mercier had already used in the same type of context in [14]. The aim of these papers was to establish controllability. The technique of exterior matrices used in [16] helped us prove controllability in a more general context than what we had done in [15]. This is why we keep this method for the present paper.

In a joint work by the authors and K. Ammari and J. Valein, the stabilization of a chain of Euler-Bernoulli beams and strings was proved using a spectral analysis also based on the exterior matrices technique (see [6]).

In this paper we will investigate the same problem as in [10]. In that paper, Chen and al. have established the exponential stability of the problem but with an assumption on the material constants (the mass densities are supposed to be decreasing and the flexural rigidities must be increasing). They remark that the assumption makes the beam more flexible at the extremity with the control and seem to think that the exponential stability could not hold without this assumption.

Our purpose is to prove, using another method (that of the exterior matrices - they have used a moment method), that in fact, the exponential stability always holds. The problem is different from the one we had studied in [16]. A feedback law is added but the interior masses have disappeared. Moreover we prove the exponential stability of the problem and not only its controllability.

The network we consider is a chain of N serially connected branches ($N \geq 2$) with $n = N + 1$ vertices (denoted by E_i).

Let us call (P_K) the stability problem:

$$\begin{aligned}
 (1) \quad & m_j u_{j,tt}(x, t) + a_j u_j^{(4)}(x, t) = 0, & \forall j \in \{1, \dots, N\}, \\
 (2) \quad & u_1(0, t) = u_1^{(1)}(0, t) = 0, \\
 (3) \quad & u_j(l_j, t) - u_{j+1}(0, t) = 0, & \forall j \in \{1, \dots, N - 1\}, \\
 (4) \quad & u_j^{(1)}(l_j, t) - u_{j+1}^{(1)}(0, t) = 0, & \forall j \in \{1, \dots, N - 1\},
 \end{aligned}$$

$$\begin{aligned}
(5) \quad & a_j u_j^{(2)}(l_j, t) - a_{j+1} u_{j+1}^{(2)}(0, t) = 0, \quad \forall j \in \{1, \dots, N-1\}, \\
(6) \quad & a_j u_j^{(3)}(l_j, t) - a_{j+1} u_{j+1}^{(3)}(0, t) = 0, \quad \forall j \in \{1, \dots, N-1\}, \\
(7) \quad & a_N u_N^{(2)}(l_N, t) = 0, \\
(8) \quad & a_N u_N^{(3)}(l_N, t) = z(t).
\end{aligned}$$

where the feedback law is chosen as :

$$(9) \quad z(t) = K u_{N,t}(l_N, t),$$

where $K > 0, t \geq 0$.

The scalar function $u_j(x, t)$ contains the information on the vertical displacement of the j -th beam. This displacement is described by the first equation where m_j is the constant mass density of the j -th beam and a_j its flexural rigidity ($1 \leq j \leq N$).

The third, fourth, fifth and sixth equations are transmission conditions. The second, seventh and eighth ones are boundary conditions. Note that the damping function $z = z(t)$ acts on the system through the exterior node E_N on the quantity $a_N u_N^{(3)}(l_N, t)$.

The goal of the paper is to establish on the first hand the existence and uniqueness of the solution of Problem (P_K) with a regularity depending on that of the initial conditions, on the other hand, that the energy of the solution decays to zero exponentially.

Before starting to study the core of the problem, we apply in Section 2 the terminology of networks to our particular network. The whole terminology can be found in early contributions of Lumer and Gramsch as well as in papers by Ali Mehmeti ([1] and [2]), von Below (cf. [8]) and Nicaise ([17] and [3]) in the eighties. We also give some properties of the spatial operator A involved in the considered evolution problem and construct an operator B and its adjoint to rewrite the problem as the abstract evolution equation (28). The aim is to apply the results of the paper by Ammari and Tucsnak ([4] and also [5]).

In Section 3, we give the main results: two properties of the eigenelements of the operator A . One called the spectral gap concerns the asymptotic behaviour of the difference between two consecutive eigenvalues, the other one is an estimate from both above and below of the value of an eigenfunction at $x = l_N$.

Both results are sufficient conditions to establish the estimate of the transfer function required by Ammari and Tucsnak's method (cf. [4] and [5]) as well as the observability inequality called (2.5) in their paper and rewritten in our context as (44). The exponential stability of the problem follows.

The proof of the spectral gap (asymptotic behaviour of the difference between two consecutive eigenvalues) is given in Section 4. This asymptotic behaviour is given by that of the roots of a function called f_∞ . In order to avoid the cancellation of the large order terms, the characteristic equation is computed using the exterior matrix method due to Paulsen (see [18]) and already used by D. Mercier in [14].

Two estimates involving the evaluation of an eigenfunction of the problem at the node where the damping acts as well as a norm of the same eigenfunction remain to be proved. It is the aim of Section 5. The choice of the basis h_i (cf. the notation at the beginning of Section 5) in which the eigenfunctions are decomposed is crucial for the asymptotic behaviour of the eigenfunctions to be studied since the expressions are very complicated especially for large values of N . In particular the exponential factor in h_3 has an important role since its presence keeps the exponential terms from being disseminated in the different matrices which would not allow an easy estimation of the involved quantities as λ tends to infinity.

2 Data and Framework

2.1 Domain and notation

The domain that we consider is a network of N ($N \in \mathbb{N} - \{0, 1\}$) serially connected beams which can be modelled by a graph $G = \bigcup_{j=1}^N k_j$. Each branch k_j having an origin and an end such that the end of the branch k_j ($1 \leq j \leq N - 1$) is connected to the beginning of the branch k_{j+1} . By the intermediary of a parametrization we will identify each branch k_j with the interval $[0, l_j]$, 0 represents the beginning of k_j and l_j the end. For each branch k_j , we fix mechanical constants $m_j > 0$ (the mass density of the beam k_j) and $a_j > 0$ (the flexural rigidity of k_j). The vibration of the branch k_j is modelled by the function $u_j(t, x)$, $t \geq 0, x \in [0, l_j]$, $j = 1, \dots, N$. The total vibration of the structure is the vectorial function $u = (u_j)_{j=1, \dots, N}$.

Notation for derivatives. In this paper, for a function $u = u(x, t)$ we make the choice to denote by u_t (u_{tt}, \dots) the first (second, ...) time derivative and $u^{(1)}$ ($u^{(2)}, \dots$) the first (second, ...) spatial derivative.

2.2 The stability problem

We assume that each beam is uniform, with constant mass density m_j and flexural rigidity a_j , $j = 1, \dots, N$. The problem, denoted by (P_K) , is the following:

$$(10) \quad m_j u_{j,tt}(x, t) + a_j u_j^{(4)}(x, t) = 0, \quad \forall j \in \{1, \dots, N\},$$

$$(11) \quad u_1(0, t) = u_1^{(1)}(0, t) = 0,$$

$$(12) \quad u_j(l_j, t) - u_{j+1}(0, t) = 0, \quad \forall j \in \{1, \dots, N - 1\},$$

$$(13) \quad u_j^{(1)}(l_j, t) - u_{j+1}^{(1)}(0, t) = 0, \quad \forall j \in \{1, \dots, N - 1\},$$

$$(14) \quad a_j u_j^{(2)}(l_j, t) - a_{j+1} u_{j+1}^{(2)}(0, t) = 0, \quad \forall j \in \{1, \dots, N - 1\},$$

$$(15) \quad a_j u_j^{(3)}(l_j, t) - a_{j+1} u_{j+1}^{(3)}(0, t) = 0, \quad \forall j \in \{1, \dots, N - 1\},$$

$$(16) \quad a_N u_N^{(2)}(l_N, t) = 0,$$

$$(17) \quad a_N u_N^{(3)}(l_N, t) = z(t).$$

The feedback law is chosen as :

$$(18) \quad z(t) = K u_{N,t}(l_N, t),$$

where $K > 0, t \geq 0$.

2.3 Abstract framework

In order to study the above system (P_K) we need to formulate it in an abstract setting. More precisely, we shall see that the framework given in [4] is well adapted to our problem (P_K) .

2.3.1 The operator A : definition and properties

For that purpose, we define the Hilbert space

$$\mathcal{H} = \prod_{j=1}^N L^2(0, l_j),$$

endowed by the inner product

$$(u, \tilde{u})_{\mathcal{H}} = \sum_{j=1}^N m_j \int_0^{l_j} \langle u_j(x), \tilde{u}_j(x) \rangle dx,$$

where $\langle \cdot, \cdot \rangle$ represents the Hermitian product in \mathbb{C} .

We also define the space V

$$V = \left\{ u \in \prod_{j=1}^N H^2(0, l_j) \text{ satisfying (19) to (21) hereafter} \right\},$$

$$(19) \quad u_1(0) = u_1^{(1)}(0) = 0,$$

$$(20) \quad u_j(l_j) - u_{j+1}(0) = 0, \quad \forall j \in \{1, \dots, N-1\},$$

$$(21) \quad u_j^{(1)}(l_j) - u_{j+1}^{(1)}(0) = 0, \quad \forall j \in \{1, \dots, N-1\},$$

as well as the sesquilinear form $a(u, v)$ for $(u, v) \in V \times V$ by

$$(22) \quad a(u, v) = \sum_{j=1}^N a_j \int_0^{l_j} \langle u_j^{(2)}(x), v_j^{(2)}(x) \rangle dx.$$

Next, we define the linear operator $A : D(A) \rightarrow \mathcal{H}$ by

$$Au = \left(\frac{a_j}{m_j} u_j^{(4)} \right)_{1 \leq j \leq N},$$

with domain

$$D(A) = \left\{ u \in \mathcal{H} : u \in \left(\prod_{j=1}^N H^4(0, l_j) \right) \cap V, \text{ satisfying (23) to (26) hereafter} \right\}$$

$$(23) \quad a_j u_j^{(2)}(l_j) - a_{j+1} u_{j+1}^{(2)}(0) = 0, \quad \forall j \in \{1, \dots, N-1\},$$

$$(24) \quad a_j u_j^{(3)}(l_j) - a_{j+1} u_{j+1}^{(3)}(0) = 0, \quad \forall j \in \{1, \dots, N-1\},$$

$$(25) \quad a_N u_N^{(2)}(l_N) = 0,$$

$$(26) \quad a_N u_N^{(3)}(l_N) = 0.$$

The operator A is a linear unbounded self-adjoint and strictly positive operator in \mathcal{H} . The domain of $A^{\frac{1}{2}}$ is $D(A^{\frac{1}{2}}) = V$ (the proof based on Friedrichs extension is left to the reader).

2.3.2 The operators B and B^*

We define the operator $B : \mathbb{C} \rightarrow V'$ such that $\forall z \in \mathbb{C}, Bz = ACz$, where V' is the dual space of V with respect to the inner product of the pivot space \mathcal{H} and $Cz = u$ is the solution of

$$u_i^{(4)} = 0, i = 1, \dots, N$$

which satisfies conditions (19), (20), (21), (23), (24), (25) as well as the following condition

$$(27) \quad a_N u_N^{(3)}(l_N) = -z.$$

Then its adjoint $B^* : V \rightarrow \mathbb{C}$ is defined by $B^*\Phi = \Phi_N(l_N)$, for any $\Phi \in V$ and the system (P_K) is described by

$$(28) \quad u_{tt}(t) + Au(t) + K \cdot BB^*u_t(t), \quad u(0) = u^0, \quad u_t(0) = u^1, \quad t \in [0, \infty).$$

with $u(t) \in V$. A solution of (28) is u such that $u(t) \in V$ for $t \in [0, \infty)$ and, for any $\Phi \in V$:

$$(29) \quad \begin{aligned} & (u_{tt}(t), \Phi)_{\mathcal{H}} + a(u(t), \Phi) + Ka(CB^*(u_t(t)), \Phi) = 0 \iff \sum_{j=1}^N m_j \int_0^{l_j} \langle u_{j,tt}(t, x), \Phi_j(x) \rangle dx \\ & + \sum_{j=1}^N a_j \int_0^{l_j} \langle u_j^{(2)}(t, x), \Phi_j^{(2)}(x) \rangle dx + K \sum_{j=1}^N a_j \int_0^{l_j} \langle p_j^{(2)}(t, x), \Phi_j^{(2)}(x) \rangle dx = 0 \end{aligned}$$

where $p = (p_1, \dots, p_N)$ is the solution of

$$p_i^{(4)} = 0, i = 1, \dots, N$$

which satisfies conditions (19), (20), (21), (23), (24), (25) and $a_N p_N^{(3)}(l_N) = -u_{N,t}(l_N, t)$.

Two integrations by parts and adapted choices for Φ lead to the equivalence between Problem (P_K) and the abstract rewriting (28). Thus, the framework given in [4] is well adapted to our problem (P_K) (see equations (1.3), (1.4) in [4]).

2.3.3 The dissipative operator \mathcal{A}_d

The dissipative operator \mathcal{A}_d introduced by Ammari and Tucsnak ([4]) in the same paper is still $\mathcal{A}_d : D(\mathcal{A}_d) \rightarrow V \times \mathcal{H}$ defined by

$$\mathcal{A}_d = \begin{pmatrix} 0 & I \\ -A & -K \cdot BB^* \end{pmatrix}$$

with domain

$$D(\mathcal{A}_d) = \{(u, v) \in V \times \mathcal{H} : Au + K \cdot BB^*v \in \mathcal{H}, v \in V\}.$$

Thus the abstract equation (28) can be rewritten as

$$(30) \quad w_t(t) = \mathcal{A}_d w(t), \quad w(0) = w^0, \quad t \in [0, \infty)$$

with $w(t) \in D(\mathcal{A}_d)$, $w^0 = (u^0, u^1)$.

Let us state some classical results based on Lumer-Phillips Theorem.

Proposition 2.1 (*existence and uniqueness of the solution, decreasing of the energy*)

1. Assume that $w^0 = (u^0, u^1) \in D(\mathcal{A}_d)$. Then equation (30) has a unique solution

$$(31) \quad w \in C(0, \infty, D(\mathcal{A}_d)) \cap C^1(0, \infty; V \times \mathcal{H}).$$

Thus equation (28) has a unique solution

$$(32) \quad u \in C^1(0, \infty; V)$$

such that $B^*u(\cdot) \in H^1(0, T; \mathbb{C})$ and

$$(33) \quad \|B^*u_t\|_{L^2(0, T; \mathbb{C})}^2 \leq C \|(u^0, u^1)\|_{V \times \mathcal{H}}^2$$

where the constant $C > 0$ is independent of (u^0, u^1) .

2. We still assume that $w^0 = (u^0, u^1) \in D(\mathcal{A}_d)$. The energy of the solution $u(t)$ given above, defined by

$$(34) \quad E(u(t)) = \frac{1}{2} \{ \|u_t(t)\|_{\mathcal{H}}^2 + \|A^{1/2}u(t)\|_{\mathcal{H}}^2 \}$$

satisfies

$$(35) \quad E(u(0)) - E(u(t)) = \int_0^t |B^*u_t(s)|^2 ds \geq 0, \quad \forall t \geq 0.$$

3 Main results

3.1 Properties of the eigenelements of the operator A

Consider the eigenvalue problem: $\lambda^2 \in \sigma(A)$ ($\lambda > 0$) is an eigenvalue of A with associated eigenvector $\phi \in D(A)$ if and only if ϕ satisfies the transmission and boundary conditions (19), (20), (21), (23), (24), (25), (26) of Section 2.3.1 and

$$(EP) \begin{cases} \phi_j^{(4)} = q_j^4 \lambda^2 \phi_j & \text{on } (0, l_j), \forall j \in \{1, \dots, N\}, \\ \phi_j \in H^4((0, l_j)), & \forall j \in \{1, \dots, N\}, \end{cases}$$

with $q_j = (m_j/a_j)^{1/4}$.

The following results are useful for stability. They are proved in Sections 4 and 5 respectively.

Theorem 3.1 (*the spectral gap*)

Let λ_k^2 , $k \in \mathbb{N}^*$, ($\lambda_k > 0$) be the (strictly) monotone increasing sequence of eigenvalues of Problem (EP) given above then

$$(36) \quad \lim_{k \rightarrow +\infty} (\lambda_{k+1} - \lambda_k) = +\infty.$$

Theorem 3.2 (*uniform estimates for $|\phi_N(l_N)|$*)

Consider the eigenvalue problem (EP) given above. For any eigenfunction $\phi \in D(A)$ associated to the eigenvalue λ^2 and for any $K > 0$, there exists two constants K_1 and K_2 such that:

$$(37) \quad K_1 \cdot \|\phi\|_{\mathcal{H}}^2 \leq |\phi_N(l_N)|^2 \leq K_2 \cdot \|\phi\|_{\mathcal{H}}^2$$

with the norm $\|\cdot\|_{\mathcal{H}}$ introduced in Section 2.3.1.

3.2 Strong stability

Using the abstract framework given in last section, we prove the decay to zero of the energy of any solution of the abstract equation (30) with suitable initial condition. It is enough to establish that the operator \mathcal{A}_d (introduced in Definition 2.3.3) has no eigenvalues on the imaginary axis.

Theorem 3.3 (*strong stability*)

It holds

$$(38) \quad \lim_{t \rightarrow +\infty} E(u(t)) = 0$$

for any solution $w = (u, v)$ of the abstract equation (30) with w^0 in $V \times \mathcal{H}$.

Proof. First we compute $\Re[(\mathcal{A}_d w, w)_{V \times \mathcal{H}}]$.

$$(39) \begin{aligned} (\mathcal{A}_d w, w)_{V \times \mathcal{H}} &= (v, u)_V + (-Au - K \cdot BB^*v, v)_{\mathcal{H}} = a(v, u) - (Au, v)_{\mathcal{H}} - K(BB^*v, v)_{\mathcal{H}} \\ &= a(u, v) - a(u, v) - K|B^*v|^2 = -2i\Im(a(u, v)) - K|v_N(l_N)|^2. \end{aligned}$$

Thus $\Re[(\mathcal{A}_d w, w)_{V \times \mathcal{H}}] = -K|v_N(l_N)|^2$.

Now we want to prove that the operator \mathcal{A}_d has no eigenvalues of the form $i\mu$, with $\mu \in \mathbb{R}$. To this end, we suppose that there exists $w = (u, v) \neq (0, 0)$ in $D(\mathcal{A}_d)$, such that $\mathcal{A}_d w = i\mu w$. Then, by definition of \mathcal{A}_d , it holds:

$$(40) \quad \begin{cases} v = i\mu u, \\ Au + K \cdot BB^*v = -i\mu v. \end{cases}$$

Now for such a w , $\Re[(\mathcal{A}_d w, w)_{V \times \mathcal{H}}] = \Re[i\mu(w, w)_{V \times \mathcal{H}}] = 0$. It follows $v_N(l_N) = 0$. Since $v_N(l_N) = B^*v$, system (40) implies $u_N(l_N) = 0$ and $Au = \mu^2 u$. Moreover, since $(u, v) \in D(\mathcal{A}_d)$, $Au + K \cdot BB^*v \in \mathcal{H}$ i.e. $Au \in \mathcal{H}$ (since $B^*v = v_N(l_N) = 0$). Thus $u \in D(A)$.

Two cases must be envisaged:

- either $\mu \neq 0$. In that case, any eigenfunction of the conservative operator A satisfies $u_N(l_N) \neq 0$ (see Theorem 3.2). This is a contradiction with our assumption.
- Or $\mu = 0$. Then, if u is an eigenfunction associated to the eigenvalue $\mu^2 = 0$,

$$0 = (Au, u)_{\mathcal{H}} = a(u, u) = \sum_{j=1}^N a_j \int_0^{l_j} \langle u_j^{(2)}(x), u_j^{(2)}(x) \rangle dx$$

which implies $u_j^{(2)} = 0$ on $(0; l_j)$ for any j in $\{0, \dots, N\}$. From condition (11), it follows $u_1 = 0$ (indeed u_1 must be a polynomial function with degree 1 which vanishes as well as its first derivative at 0). Then conditions (12) and (13) combined with $u_1(l_1) = u_2(0) = 0$ and $u_2^{(2)} = 0$ on $(0; l_2)$ imply $u_2 = 0$ and so on: $u_j = 0$ on $(0; l_j)$ for any j in $\{0, \dots, N\}$. This is a contradiction.

The conclusion is that \mathcal{A}_d has no eigenvalues of the form $i\mu$, with $\mu \in \mathbb{R}$. The result follows, using the main theorem of [7]. ■

3.3 Exponential stability

Using Theorem 2.2 of [4] we can state the exponential stability of our system. To this end, we prove (1.5) of [4] (also called hypothesis (H) , which is the estimate of a transfer function) directly (using the orthonormal basis formed by the eigenfunctions of the operator A) as well as the estimate (2.5) which is an observability inequality. To establish (H) , we need the spectral gap and since the proof of the spectral gap is a long and technical proof based on the same ideas as those we have already used in a previous paper (cf. [16]), it is given in next section. As for the observability inequality (2.5), its proof requires both the spectral gap and a uniform estimate for $|\phi_N(l_N)|$ which is proved via an adaptation of ancient results (same paper [16]). It is also given in another section. Note that some new technical difficulties appear in the

calculations and proofs of Sections 4 and 5 and that, although they are based on the same ideas as a previous paper, they are not trivial at all.

Theorem 3.4 (*exponential stability*)

The system described by the abstract equation (30) is exponentially stable in $V \times \mathcal{H}$.

Proof. To prove this theorem, we use Theorem 2.2 of [4]. In the **first part of the proof**, we will check that the following hypothesis (H) is satisfied:

(H): If $\beta > 0$ is fixed and $C_\beta = \{\lambda \in \mathbb{C} | \Re(\lambda) = \beta\}$, the function

$$\lambda \in C_\beta \rightarrow H(\lambda) = \lambda B^*(\lambda^2 + A)^{-1}B \in \mathcal{L}(\mathbb{C})$$

is bounded.

Since the proof is long, we divide it into several steps.

First step: rewrite $H(\lambda)$ as a series. We start by computing $B(1)$ (B is a linear operator defined on \mathbb{C} thus computing $B(1)$ is enough to know any value $B(\lambda)$).

Since $B(1) \in V'$ then there exists a sequence $(\alpha_k)_{k \in \mathbb{N}^*}$ such that

$$\sum_{k=1}^{\infty} \left| \frac{\alpha_k}{\lambda_k} \right|^2 < \infty \text{ and } B(1) = \sum_{k=1}^{\infty} \alpha_k \phi_k.$$

(with V introduced in Definition 2.3.1, $(\phi_k)_k$ the orthonormal basis formed by the eigenvectors of the operator A and λ_k^2 the eigenvalues).

Let $h = \sum_{k=1}^{\infty} h_k \phi_k$ any element of V (i.e $\sum_{k=1}^{\infty} |h_k \lambda_k|^2 < \infty$) then

$$(41) \quad \langle B(1), h \rangle_{V',V} = \sum_{k=1}^{\infty} \langle \alpha_k, h_k \rangle .$$

We also have

$$(42) \quad \langle B(1), h \rangle_{V',V} = \langle 1, B^*h \rangle_{\mathbb{C},\mathbb{C}} = \overline{h_N(l_N)} = \sum_{k=1}^{\infty} \overline{h_k} \overline{\phi_{k,N}(l_N)}.$$

From (41) and (42) we deduce that

$$B(1) = \sum_{k=1}^{\infty} \overline{\phi_{k,N}(l_N)} \phi_k.$$

Now, as previously, we directly compute $(\lambda^2 I + A)^{-1}B(1)$ in the orthonormal basis $(\phi_k)_k$ and easily find that

$$(\lambda^2 I + A)^{-1}B(1) = \sum_{k=1}^{\infty} \frac{\overline{\phi_{k,N}(l_N)}}{\lambda^2 + \lambda_k^2} \phi_k.$$

Therefore

$$H(\lambda) = \lambda B^*(\lambda^2 I + A)^{-1}B(1) = \lambda \sum_{k=1}^{\infty} \frac{|\phi_{k,N}(l_N)|^2}{\lambda^2 + \lambda_k^2}.$$

Second step: find an estimate for the imaginary part of $H(\lambda)/\lambda$ on the line C_β . We can choose $\beta = 1$ without loss of generality. Calculating the imaginary part of $H(1+iy)/(1+iy)$ leads to:

$$-\frac{1}{2y} \Im \left(\frac{H(\lambda)}{\lambda} \right) = \sum_{k=1}^{\infty} |\phi_{k,N}(l_N)|^2 \cdot f_1(y, \lambda_k), \text{ for } \lambda = 1 + iy, y \in \mathbb{R}.$$

where $f_1(y, \lambda) = \frac{1}{4y^2 + (1 - y^2 + \lambda^2)^2}$.

Due to Theorem 3.2, $K_1 \leq |\phi_{k,N}(l_N)|^2 \leq K_2$ (the norm of each ϕ_k in \mathcal{H} is one).

Thus, the function we need to estimate from above is the function of y defined by

$$\Sigma(y) = \sum_{k=1}^{\infty} \frac{1}{4y^2 + (1 - y^2 + \lambda^2)^2}.$$

For a fixed $y > 0$, the sum Σ is separated into three terms: one with the small values of λ_k , one with the values of λ_k which are close to the value of y (these values contribute mostly to the sum but their number is finite) and the last one with the big values of λ_k .

1. By definition, $\Sigma_1(y) = \sum_{k:\lambda_k \leq y - \sqrt{y}} \frac{1}{4y^2 + (1 - y^2 + \lambda^2)^2}$. The function f_1 is an increasing function of λ as soon as y is large enough. Indeed the derivative $\partial_\lambda f_1(y, \lambda)$ is equal to

$$\frac{4\lambda(y^2 - \lambda^2 - 1)}{(4y^2 + (1 - y^2 + \lambda^2)^2)^2}.$$

Then, since $\lambda \leq y - \sqrt{y}$, $f_1(y, \lambda) \leq f_1(y, y - \sqrt{y})$ and

$$f_1(y, y - \sqrt{y}) = \frac{1}{4y^2 + (1 - y^2 + (y - \sqrt{y})^2)^2} \lesssim \frac{1}{y^4}.$$

Note that the notation $A \lesssim B$ means the existence of a positive constant C , which is independent of A and B such that $A \leq CB$.

Now, from the spectral gap (cf. Theorem 3.1), $\lambda_k \gtrsim k^2$. Then, denoting by \mathcal{N}_y , the number of λ_k such that $\lambda_k \leq y - \sqrt{y}$, it holds:

$$(\mathcal{N}_y)^2 \lesssim \lambda_{\mathcal{N}_y} \leq y - \sqrt{y}.$$

Thus, for large values of y , $\Sigma_1(y) \lesssim \mathcal{N}_y \frac{1}{y^4} \lesssim \frac{\sqrt{y}}{y^4} \lesssim \frac{1}{y^2}$.

2. By definition, $\Sigma_2(y) = \sum_{k:y - \sqrt{y} \leq \lambda_k \leq y + \sqrt{y}} \frac{1}{4y^2 + (1 - y^2 + \lambda^2)^2}$.

First $f_1(y, \lambda) \leq \frac{1}{4y^2}$.

Secondly, the number N_y which is the number of λ_k such that $y - \sqrt{y} \leq \lambda_k \leq y + \sqrt{y}$ is bounded as regards y . Indeed N_y is smaller than the quotient of the amplitude of the interval $(2\sqrt{y})$ over the minimum gap between two consecutive values of λ_k (with $y - \sqrt{y} \leq \lambda_k \leq y + \sqrt{y}$). Now the gap between two values of λ_k is:

$$(43) \quad \begin{aligned} \lambda_{k+1} - \lambda_k &= (\sqrt{\lambda_{k+1}} - \sqrt{\lambda_k})(\sqrt{\lambda_{k+1}} + \sqrt{\lambda_k}) \\ &\geq \sigma(\sqrt{\lambda_{k+1}} + \sqrt{\lambda_k}) \geq 2\sigma\sqrt{\lambda_k} \gtrsim k \end{aligned}$$

where the parameter σ is introduced in the proof of Theorem 4.10.

Now, we define k_0 the largest integer such that $\lambda_{k_0} \leq y - \sqrt{y}$. Then, for any k such that $y - \sqrt{y} \leq \lambda_k \leq y + \sqrt{y}$, $\lambda_{k+1} - \lambda_k \gtrsim k_0$. Thus $N_y \lesssim \frac{2\sqrt{y}}{k_0}$.

Moreover there exists $\gamma_2 > 0$, $\lambda_k \leq \gamma_2 k^2$ (cf. [11]) and since $y - \sqrt{y} \leq \lambda_{k_0+1} \leq \gamma_2(k_0+1)^2$, $k_0 \geq \sqrt{\frac{y - \sqrt{y}}{\gamma_2}} - 1$ and $N_y \lesssim \frac{2\sqrt{y}}{\sqrt{y}} \lesssim 1$.

The conclusion is: $\Sigma_2(y) \lesssim \frac{1}{y^2}$.

3. By definition, $\Sigma_3(y) = \sum_{k:y+\sqrt{y} \leq \lambda_k} \frac{1}{4y^2 + (1 - y^2 + \lambda^2)^2}$. The derivative $\partial_y f_1(y, \lambda)$ is equal to

$$\frac{4y(\lambda^2 - y^2 - 1)}{(4y^2 + (1 - y^2 + \lambda^2)^2)^2}.$$

If $\lambda \geq y + \sqrt{y}$, it is positive then $f_1(y, \lambda) \leq f_1(y(\lambda), \lambda)$ with $y(\lambda) + \sqrt{y(\lambda)} = \lambda$ and $y(\lambda) \leq \lambda$ i.e. $y(\lambda) = \frac{1}{2}(2\lambda + 1 - \sqrt{4\lambda + 1})$. Since $\lambda_k \gtrsim k^2$ (as it was said above), the series $\sum_k \frac{1}{\lambda_k}$ is convergent and it follows, after calculations:

$$f_1(y, \lambda) \lesssim \frac{1}{\lambda^3} \text{ which implies } \Sigma_3(y) \lesssim \frac{1}{y^2} \sum_{k:y+\sqrt{y} \leq \lambda_k} \frac{1}{\lambda_k} \lesssim \frac{1}{y^2}.$$

The conclusion of this part is that the imaginary part of $H(\lambda)/\lambda$ satisfies:

$$\left| \Im \left(\frac{H(\lambda)}{\lambda} \right) \right| \lesssim \frac{1}{y}, \text{ for any } \lambda \in C_\beta = \{\lambda \in \mathbb{C} | \Re(\lambda) = \beta\}.$$

Third step: find an estimate for the real part of $H(\lambda)/\lambda$ on the line C_β . We still choose $\beta = 1$ and calculate the real part of $H(1 + iy)/(1 + iy)$. It is:

$$\Re \left(\frac{H(\lambda)}{\lambda} \right) = \sum_{k=1}^{\infty} |\phi_{k,N}(l_N)|^2 \cdot f_2(y, \lambda_k), \text{ for } \lambda = 1 + iy, y \in \mathbb{R}.$$

where $f_2(y, \lambda) = \frac{1 - y^2 + \lambda^2}{4y^2 + (1 - y^2 + \lambda^2)^2}$. We still separate the sum into three terms denoted by Σ'_1 , Σ'_2 and Σ'_3 .

Since the ideas of the proof are similar to those of the proof for the imaginary part, we give less details for this step.

For the estimate of Σ'_1 , we first prove that the derivative $\partial_\lambda f_2(y, \lambda)$ is negative if $\lambda \leq y - \sqrt{y}$ and y is large enough. Thus $|f_2(y, \lambda)| = -f_2(y, \lambda) \leq -f_2(y, y - \sqrt{y}) \lesssim \frac{1}{y^{3/2}}$.

Now the number \mathcal{N}_y of λ_k such that $\lambda_k \leq y - \sqrt{y}$ satisfies $\mathcal{N}_y \lesssim \sqrt{y}$ since $(\mathcal{N}_y)^2 \lesssim \lambda_{\mathcal{N}_y} \leq y - \sqrt{y}$. Thus $\Sigma_1(y) \lesssim \frac{1}{y}$.

For the estimate of Σ'_2 , we prove that the derivative $\partial_\lambda f_2(y, \lambda)$ vanishes at λ such that $1 + \lambda^2 - y^2 = \pm 2y$ i.e. at $\lambda_\pm = \sqrt{y^2 \pm 2y} - 1$. Moreover, $f_2(y, y \pm \sqrt{y}) = \frac{\pm 1}{2y\sqrt{y}} + o\left(\frac{\pm 1}{2y\sqrt{y}}\right)$. Since $f_2(y, \lambda_\pm) = \frac{\pm 1}{4y}$, it holds, for any λ in $[y - \sqrt{y}; y + \sqrt{y}]$, $|f_2(y, \lambda)| \leq |f_2(y, \lambda_+)| \lesssim \frac{1}{y}$. Thus $\Sigma'_2 \lesssim \frac{1}{y}$ (recall that N_y is bounded, cf. the second step).

At last, for the estimate of Σ'_3 , we prove that the derivative $\partial_y f_2(y, \lambda)$ is positive if $\lambda \geq y + \sqrt{y}$ and y is large enough. Thus, $f_2(y, \lambda) \leq f_2(\lambda, \lambda)$ and since $f_2(\lambda, \lambda) = \frac{1}{4\lambda^2 + 1} \lesssim \frac{1}{\lambda^2}$, it follows:

$$\Sigma'_3(y) \lesssim \frac{1}{y} \sum_{k: y + \sqrt{y} \leq \lambda_k} \frac{1}{\lambda_k} \lesssim \frac{1}{y}.$$

The conclusion of this part is that the real part of $H(\lambda)/\lambda$ satisfies:

$$\left| \Re \left(\frac{H(\lambda)}{\lambda} \right) \right| \lesssim \frac{1}{y}, \text{ for any } \lambda \in C_\beta = \{\lambda \in \mathbb{C} | \Re(\lambda) = \beta\}.$$

The **second assumption** for the exponential stability is the observability inequality called (2.5) in the paper by Ammari and Tucsnak. It follows from the spectral gap and the uniform estimate given in Section 3.1 using a result due to Haraux (cf. [12]). Indeed (2.5) is:

$$(44) \quad \exists T > 0, \exists C(T), \int_0^T |v_N(l_N, t)|^2 dt \geq C(T) \cdot \|U_0\|_{V \times H}^2$$

where $U(t) = (u(t), v(t))^t$ satisfies $U' = \mathcal{A}_c U$, $U(0) = U_0 \in D(\mathcal{A}_c)$, with \mathcal{A}_c the conservative operator defined like \mathcal{A}_d with $K = 0$.

Define $\Phi_k = (\phi_k, i\lambda_k \phi_k)^t$ an orthonormal basis of eigenfunctions of the operator \mathcal{A}_c (in particular $\|\Phi_k\|_{V \times H} = 1$). The result due to Haraux (cf. [12]) that we already used with more details in [16] allows to write:

$$(45) \quad \exists T > 0, \exists C_1(T), \int_0^T |v_N(l_N, t)|^2 dt \geq C_1(T) \sum_{k=1}^{\infty} |u_0^k|^2 |\lambda_k|^2 |\phi_{k,N}(l_N)|^2$$

where $U_0 = \sum_{k=1}^{\infty} u_0^k \Phi_k$. It follows from Theorem 3.2, $|\phi_{k,N}(l_N)|^2 \geq K_1 \|\phi_k\|_H^2$ and since there exists C_2 such that $\|\phi_k\|_H^2 \geq C_2 \frac{1}{|\lambda_k|^2} \|\Phi_k\|_{V \times H}^2$

$$(46) \quad \exists T > 0, \exists C_1(T), \exists C_2, \exists K_1, \int_0^T |v_N(l_N, t)|^2 dt \geq C_1(T) C_2 K_1 \sum_{k=1}^{\infty} |u_0^k|^2.$$

Assumption (44) follows with $C(T) = C_1(T) C_2 K_1$ since $\|U_0\|_{V \times H}^2 = \sum_{k=1}^{\infty} |u_0^k|^2$. ■

4 Proof of the spectral gap using exterior matrices

The proof of the spectral gap follows the lines of the proof of our previous paper on the boundary controllability of a chain of serially connected Euler-Bernoulli beams with interior masses (cf. [16]). We also use the exterior matrix method due to W. H. Paulsen (see [18]).

We need to determine the asymptotic behaviour of the characteristic equation of the eigenvalue problem: $\lambda^2 \in \sigma(A)$ ($\lambda > 0$) is an eigenvalue of A with associated eigenvector $\phi \in D(A)$ if and only if ϕ satisfies the transmission and boundary conditions (19), (20), (21), (23), (24), (25), (26) of Section 2.3.1 and

$$(EP) \begin{cases} \phi_j^{(4)} = q_j^4 \lambda^2 \phi_j & \text{on } (0, l_j), \forall j \in \{1, \dots, N\}, \\ \phi_j \in H^4((0, l_j)), & \forall j \in \{1, \dots, N\}, \end{cases}$$

with $q_j = (m_j/a_j)^{1/4}$.

4.1 Recall of notation and of some properties

Let ϕ be a non-trivial solution of the above eigenvalue problem (EP) and λ^2 ($\lambda > 0$) be the corresponding eigenvalue.

For each $j \in \{1, \dots, N\}$, the vector function V_j is defined by

$$V_j(x) = \left(\phi_j(x), \frac{\phi_j^{(1)}(x)}{\sqrt{\lambda}}, a_j \frac{\phi_j^{(2)}(x)}{\lambda}, a_j \frac{\phi_j^{(3)}(x)}{\lambda \sqrt{\lambda}} \right)^t, \forall x \in [0, l_j].$$

Keeping the notation a_j and l_j introduced in Section 2, the matrix A_j is $A_j := A(q_j, b_j, m_j)$ with $q_j = (m_j/a_j)^{1/4}$, $b_j = q_j l_j$ and $A(q, b, m)$ the square matrix of order 4 defined by

$$(47) \quad A(q, b, m) = \frac{1}{2} \begin{pmatrix} ch + c & \frac{sh + s}{q} & \frac{q^2(ch - c)}{m} & \frac{q(sh - s)}{m} \\ q(sh - s) & ch + c & \frac{q^3(sh + s)}{m} & \frac{q^2(ch - c)}{m} \\ \frac{m(ch - c)}{q^2} & \frac{m(sh - s)}{q^3} & ch + c & \frac{m}{sh + s} \\ \frac{m}{q} & \frac{m}{q^2} & q(sh - s) & ch + c \end{pmatrix}$$

with the notation $c = \cos(b\sqrt{\lambda})$, $s = \sin(b\sqrt{\lambda})$, $ch = \cosh(b\sqrt{\lambda})$, $sh = \sinh(b\sqrt{\lambda})$. To finish with, the matrix $M(\lambda)$ is given by

$$(48) \quad M(\lambda) = A_N A_{N-1} \dots A_2 A_1.$$

Lemma 4.1 *(a few trivial but useful properties)*

With the notation introduced above:

$$\begin{aligned} V_j(l_j) &= A_j V_j(0), \quad \forall j \in \{1, \dots, N\}, \\ V_{j+1}(0) &= V_j(l_j), \quad \forall j \in \{1, \dots, N-1\}, \\ V_N(l_N) &= M(\lambda) V_1(0). \end{aligned}$$

The proof is analogous to that of [16].

Theorem 4.2 *(the characteristic equation for the eigenvalue problem corresponding to a chain of N branches)*

$\lambda^2 > 0$ is an eigenvalue of A if and only if λ satisfies the characteristic equation

$$(49) \quad f(\sqrt{\lambda}) = \det(M_{22}(\lambda)) = 0,$$

where $M_{22}(\lambda)$ is the square matrix of order 2 which is the restriction of the matrix $M(\lambda)$, given by (48), to its last two lines and its last two columns.

For that property again, the proof is similar to that of [16].

4.2 Rewriting of the characteristic equation using the exterior matrix method

The exterior matrix method presented in [18] is a very useful method which allows to compute asymptotically the eigenfrequencies for the vibrations of serially connected elements which are governed by fourth-order equations. But our goal is to get the spectral gap. The main idea is to exploit the special properties of the exterior matrices associated to our problem in order to obtain the desired results.

The whole section makes use of the same ideas as in a previous paper by D. Mercier (see [14]).

First, we simply recall the definition of exterior matrix and some useful results that we need in the sequel (see [18] for more details).

Definition 4.3 If $M = (m_{ij})$ is a 4×4 matrix, then the exterior matrix of M is the 6×6 matrix given by:

$$\text{ext}(M) = \begin{pmatrix} \text{ext}(M)_{11} & \text{ext}(M)_{12} \\ \text{ext}(M)_{21} & \text{ext}(M)_{22} \end{pmatrix},$$

where each block $\text{ext}(M)_{ij}$, $i, j = 1, 2$, is a 3×3 matrix given hereafter:

$$\text{ext}(M)_{11} = \begin{pmatrix} \left| \begin{array}{cc} m_{11} & m_{12} \\ m_{21} & m_{22} \end{array} \right| & \left| \begin{array}{cc} m_{13} & m_{12} \\ m_{21} & m_{23} \end{array} \right| & \left| \begin{array}{cc} m_{11} & m_{14} \\ m_{21} & m_{24} \end{array} \right| \\ \left| \begin{array}{cc} m_{11} & m_{12} \\ m_{31} & m_{32} \end{array} \right| & \left| \begin{array}{cc} m_{11} & m_{13} \\ m_{31} & m_{33} \end{array} \right| & \left| \begin{array}{cc} m_{11} & m_{14} \\ m_{31} & m_{34} \end{array} \right| \\ \left| \begin{array}{cc} m_{11} & m_{12} \\ m_{41} & m_{42} \end{array} \right| & \left| \begin{array}{cc} m_{11} & m_{13} \\ m_{41} & m_{43} \end{array} \right| & \left| \begin{array}{cc} m_{11} & m_{14} \\ m_{41} & m_{44} \end{array} \right| \end{pmatrix},$$

$$\text{ext}(M)_{12} = \begin{pmatrix} \left| \begin{array}{cc} m_{13} & m_{14} \\ m_{23} & m_{24} \end{array} \right| & - \left| \begin{array}{cc} m_{12} & m_{14} \\ m_{22} & m_{24} \end{array} \right| & \left| \begin{array}{cc} m_{12} & m_{13} \\ m_{22} & m_{23} \end{array} \right| \\ \left| \begin{array}{cc} m_{13} & m_{14} \\ m_{33} & m_{34} \end{array} \right| & - \left| \begin{array}{cc} m_{12} & m_{14} \\ m_{32} & m_{34} \end{array} \right| & \left| \begin{array}{cc} m_{12} & m_{13} \\ m_{32} & m_{33} \end{array} \right| \\ \left| \begin{array}{cc} m_{13} & m_{14} \\ m_{43} & m_{44} \end{array} \right| & - \left| \begin{array}{cc} m_{12} & m_{14} \\ m_{42} & m_{44} \end{array} \right| & \left| \begin{array}{cc} m_{12} & m_{13} \\ m_{42} & m_{43} \end{array} \right| \end{pmatrix},$$

$$\text{ext}(M)_{21} = \begin{pmatrix} \left| \begin{array}{cc} m_{31} & m_{32} \\ m_{41} & m_{42} \end{array} \right| & \left| \begin{array}{cc} m_{31} & m_{33} \\ m_{41} & m_{43} \end{array} \right| & \left| \begin{array}{cc} m_{31} & m_{34} \\ m_{41} & m_{44} \end{array} \right| \\ - \left| \begin{array}{cc} m_{21} & m_{22} \\ m_{41} & m_{42} \end{array} \right| & - \left| \begin{array}{cc} m_{21} & m_{23} \\ m_{41} & m_{43} \end{array} \right| & - \left| \begin{array}{cc} m_{21} & m_{24} \\ m_{41} & m_{44} \end{array} \right| \\ \left| \begin{array}{cc} m_{21} & m_{22} \\ m_{31} & m_{32} \end{array} \right| & \left| \begin{array}{cc} m_{21} & m_{23} \\ m_{31} & m_{33} \end{array} \right| & \left| \begin{array}{cc} m_{21} & m_{24} \\ m_{31} & m_{34} \end{array} \right| \end{pmatrix},$$

$$ext(M)_{22} = \begin{pmatrix} \begin{vmatrix} m_{33} & m_{34} \\ m_{43} & m_{44} \end{vmatrix} & - \begin{vmatrix} m_{32} & m_{34} \\ m_{42} & m_{44} \end{vmatrix} & \begin{vmatrix} m_{32} & m_{33} \\ m_{42} & m_{43} \end{vmatrix} \\ - \begin{vmatrix} m_{23} & m_{24} \\ m_{43} & m_{44} \end{vmatrix} & \begin{vmatrix} m_{22} & m_{24} \\ m_{42} & m_{44} \end{vmatrix} & - \begin{vmatrix} m_{22} & m_{23} \\ m_{42} & m_{43} \end{vmatrix} \\ \begin{vmatrix} m_{23} & m_{24} \\ m_{33} & m_{34} \end{vmatrix} & - \begin{vmatrix} m_{32} & m_{34} \\ m_{22} & m_{24} \end{vmatrix} & \begin{vmatrix} m_{22} & m_{23} \\ m_{32} & m_{33} \end{vmatrix} \end{pmatrix}.$$

Lemma 4.4 *If M_1 and M_2 are 4×4 matrices, then*

$$ext(M_1 M_2) = ext(M_1) ext(M_2).$$

Proof. Sketch of the proof (for more details see Lemma 1 of [18].)

Given a matrix $M \in \mathcal{M}_4(\mathbb{R})$, we define a linear map M^* in $\mathcal{M}_4(\mathbb{R})$ such that :

$$(50) \quad \forall A \in \mathcal{M}_4(\mathbb{R}), M^*(A) = MAM^T.$$

It is easy to prove that the map $M \rightarrow M^*$ is a homomorphism (i.e we have $M_1^* M_2^* = (M_1 M_2)^*$) and that M^* sends anti-symmetric matrices to anti-symmetric matrices, so we can restrict M^* to this subspace. A basis for the 4×4 anti-symmetric matrices is

$$e_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, e_2 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, e_3 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix},$$

$$e_4 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}, e_5 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, e_6 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Using this basis, we find that M^* , when restricted to anti-symmetric matrices can be expressed by the 6×6 matrix $ext(M)$ given in Definition 4.3.

(50) expresses that the map $M \rightarrow M^*$ is a homomorphism. ■

Theorem 4.5 *(the characteristic equation rewritten in terms of exterior matrices)*

Let $\lambda^2 > 0$ be an eigenvalue of A then λ satisfies the characteristic equation

$$(51) \quad f(\sqrt{\lambda}) = e_4^t ext(M(\lambda)) e_4 = 0,$$

or equivalently

$$(52) \quad f(\sqrt{\lambda}) = e_4^t ext(A_N) ext(A_{N-1}) \dots ext(A_1) e_4 = 0,$$

where $M(\lambda)$ is the square matrix of order 4 given by (48) and $e_4^t = (0, 0, 0, 1, 0, 0)$.

Proof. The proof is analogous to that of the paper [16]. The only difference is that the determinant of M_{22} is the term of the 4-th line and 4-th column of the matrix $ext(M)$. ■

4.3 The asymptotic behaviour of the characteristic equation

As in [14], we study the asymptotic behaviour of the exterior matrices involved in $\text{ext}(M(\lambda))$ in order to get the asymptotic behaviour of the characteristic equation (52) as λ tends to ∞ . This is enough to establish the following property called spectral gap. Let λ_k^2 , $k \in \mathbb{N}^*$, ($\lambda_k > 0$) be the (strictly) monotone increasing sequence of eigenvalues of Problem (EP) given at the beginning of Section 4 then

$$(53) \quad \lim_{k \rightarrow +\infty} (\lambda_{k+1} - \lambda_k) = +\infty.$$

Thus the aim of the following is to study the asymptotic behaviour of the exterior matrix of each matrix A_j contained in the expression (48). From now on the notation $o(\lambda)$ is used for a square matrix of the appropriate size such that all its terms are dominated by the function $\lambda \mapsto \lambda$ asymptotically.

Definition 4.6 (definition of the matrices C , S and of the vectors V_1 and V_2)

$$(54) \quad C(q, m) = \begin{pmatrix} 1 & \frac{q^3}{m} & 0 & -\frac{q^4}{m^2} & \frac{q}{m} & 0 \\ \frac{m}{q^3} & 2 & \frac{1}{q} & -\frac{q}{m} & 0 & \frac{1}{q} \\ 0 & q & 1 & 0 & -\frac{1}{q} & 1 \\ -\frac{m^2}{q^4} & -\frac{m}{q} & 0 & 1 & -\frac{m}{q^3} & 0 \\ \frac{m}{q} & 0 & -q & -\frac{q^3}{m} & 2 & -q \\ 0 & q & 1 & 0 & -\frac{1}{q} & 1 \end{pmatrix}.$$

$$(55) \quad S(q, m) = \begin{pmatrix} 0 & \frac{q^3}{m} & \frac{q^2}{m} & 0 & -\frac{q}{m} & \frac{q^2}{m} \\ -\frac{m}{q^3} & 0 & \frac{1}{q} & \frac{q}{m} & -\frac{m}{q^2} & \frac{1}{q} \\ -\frac{m}{q^2} & -q & 0 & \frac{q^2}{m} & -\frac{1}{q} & 0 \\ 0 & -\frac{m}{q} & -\frac{m}{q^2} & 0 & \frac{m}{q^3} & -\frac{m}{q^2} \\ \frac{m}{q} & 2q^2 & q & -\frac{q^3}{m} & 0 & q \\ -\frac{m}{q^2} & -q & 0 & \frac{q^2}{m} & -\frac{1}{q} & 0 \end{pmatrix}.$$

$$(56) \quad V_1(q, m) = \left(-\frac{q^4}{m^2}, -\frac{q}{m}, 0, 1, -\frac{q^3}{m}, 0 \right)^t.$$

$$(57) \quad V_2(q, m) = \left(0, \frac{q}{m}, \frac{q^2}{m}, 0, -\frac{q^3}{m}, \frac{q^2}{m} \right)^t.$$

Lemma 4.7 (properties of H , C , S , V_1 and V_2)

The exterior matrix of A_j may be written as:

$$\text{ext}(A_j) = e^{b_j\sqrt{\lambda}}H(q_j, b_j, m_j) + o(e^{b_j\sqrt{\lambda}}).$$

where the matrix H is:

$$(58) \quad H(q_j, b_j, m_j) = \cos(b_j\sqrt{\lambda})C(q_j, m_j) + \sin(b_j\sqrt{\lambda})S(q_j, m_j).$$

The rank of both matrices $C(q, m)$ and $S(q, m)$ defined above is 2. A basis for the range of $C(q, m)$ is $\{V_1(q, m), V_2(q, m)\}$. Idem for the range of $S(q, m)$.

Proof. The first property is proved via a long calculation using the definition of exterior matrices. We do not give the details here. But for instance the first coefficient of $\text{ext}(A_j)$ is :

$$\frac{1}{2} \left(1 + \cos(b_j\sqrt{\lambda}) \cdot \cosh(b_j\sqrt{\lambda}) \right) = \frac{e^{b_j\sqrt{\lambda}}}{4} \cos(b_j\sqrt{\lambda}) + o(e^{b_j\sqrt{\lambda}}).$$

A simple computation for the matrix $C(q, m)$ leads to $L_3 = q \cdot \left(\frac{q^3}{m}L_2 - L_1 \right)$, $L_4 = -\frac{m^2}{q^4}L_1$, $L_5 = \left(q^2 + \frac{m}{q} \right) L_1 - \frac{q^5}{m}L_2$ and $L_6 = L_3$. And for the matrix $S(q, m)$, $L_2 = \frac{2q^3}{m}L_1 - \frac{1}{q^2}L_5$, $L_3 = \frac{q^4}{m}L_1 - \frac{1}{q}L_5$, $L_4 = \frac{1}{q^4}L_1$ and $L_6 = L_3$. Hence the rank is 2 for both matrices.

The vector V_1 is in the range of C and S since $V_1 = C(0, 0, 0, 1, 0, 0)^t$ and $V_1 = S(0, 0, 0, 1, q^3/m, 0)^t$. Likewise for V_2 : $V_2 = C(0, 0, 0, -1, -q^3/m, 0)^t$ and $V_2 = S(0, 0, 0, 1, 0, 0)^t$.

At last V_1 and V_2 satisfy: $\lambda_1 V_1(q, m) + \lambda_2 V_2(q, m) = 0$ implies $\lambda_1 = \lambda_2 = 0$. ■

Lemma 4.8 For all $j = 1, \dots, N - 1$ we have

$$(59) \quad \left(\prod_{i=j}^1 H(q_i, b_i, m_i) \right) \cdot e_4 = \alpha_j(\sqrt{\lambda})V_1(q_j, m_j) + \beta_j(\sqrt{\lambda})V_2(q_j, m_j)$$

where $V_k(q, m)$, $k = 1, 2$ are introduced in Definition 4.6 and e_4 in Theorem 4.5, $\alpha_j(\cdot), \beta_j(\cdot)$ are trigonometrical polynomials which only depend on $q_i, b_i, m_i, i = 1, \dots, j$.

Moreover, there exists a constant $d_j > 0$ (which only depends on the material constants) such that the Wronskian $W_j(x) = \alpha_j(x)\beta_j'(x) - \alpha_j'(x)\beta_j(x)$ satisfies

$$(60) \quad W_j(x) \geq d_j > 0, \forall x \in \mathbb{R}.$$

Proof. We argue by iteration. We suppose that $j = 1$. By Definition 4.6 and Lemma 4.7, it holds:

$$H(q_1, b_1, m_1)e_4 = \cos(b_1\sqrt{\lambda})V_1(q_1, m_1) + \sin(b_1\sqrt{\lambda})V_2(q_1, m_1).$$

Thus (59) holds for $j = 1$ with $\alpha_1(x) = \cos(b_1x)$, $\beta_1(x) = \sin(b_1x)$. Since $\forall x \in \mathbb{R}$, $W_1(x) = b_1$, then, for $j = 1$, (60) is true with $d_1 = b_1 > 0$.

Now, suppose that (59) holds for $j - 1$ and that there exists a constant d_{j-1} such that:

$\forall x \in \mathbb{R}, W_{j-1}(x) \geq d_{j-1} > 0.$

Thus, with (58) of Lemma 4.7 we may write:

$$\begin{aligned}
(61) \quad \left(\prod_{i=j}^1 H(q_i, b_i, m_i) \right) e_4 &= H(q_j, b_j, m_j) (\alpha_{j-1}(\sqrt{\lambda}) V_1(q_{j-1}, m_{j-1}) \\
&+ \beta_{j-1}(\sqrt{\lambda}) V_2(q_{j-1}, m_{j-1})) \\
&= (\cos(b_j \sqrt{\lambda}) C(q_j, m_j) + \sin(b_j \sqrt{\lambda}) S(q_j, m_j)) \\
&\times (\alpha_{j-1}(\sqrt{\lambda}) V_1(q_{j-1}, m_{j-1}) + \beta_{j-1}(\sqrt{\lambda}) V_2(q_{j-1}, m_{j-1})).
\end{aligned}$$

Now, from Lemma 4.7, we know that there exist constants $z_i, i = 1, \dots, 8.$ such that

$$\begin{aligned}
(62) \quad C(q_j, m_j) V_1(q_{j-1}, m_{j-1}) &= z_1 V_1(q_j, m_j) + z_2 V_2(q_j, m_j), \\
C(q_j, m_j) V_2(q_{j-1}, m_{j-1}) &= z_3 V_1(q_j, m_j) + z_4 V_2(q_j, m_j), \\
S(q_j, m_j) V_1(q_{j-1}, m_{j-1}) &= z_5 V_1(q_j, m_j) + z_6 V_2(q_j, m_j), \\
S(q_j, m_j) V_2(q_{j-1}, m_{j-1}) &= z_7 V_1(q_j, m_j) + z_8 V_2(q_j, m_j).
\end{aligned}$$

Using the expressions of $C(q_j, m_j), S(q_j, m_j), V_1(q_{j-1}, m_{j-1})$ and $V_2(q_{j-1}, m_{j-1})$ given in Definition 4.6 we get after some computations:

$$\begin{aligned}
(63) \quad z_1 = z_6 &= \frac{(m_{j-1} q_j + m_j q_{j-1})(m_{j-1} q_j^3 + m_j q_{j-1}^3)}{m_{j-1} q_j^3}, \\
z_2 = z_3 = -z_5 = z_8 &= \frac{m_j q_{j-1} (q_j^2 - q_{j-1}^2)}{m_{j-1} q_j^3}, \\
z_4 = -z_7 &= \frac{m_j q_{j-1} (q_j + q_{j-1})^2}{m_{j-1} q_j^3}.
\end{aligned}$$

Using (62) in the development of the last expression of (61)

$$\left(\prod_{i=j}^1 H(q_i, b_i, m_i) \right) e_4 = \alpha_j(\sqrt{\lambda}) V_1(q_j, m_j) + \beta_j(\sqrt{\lambda}) V_2(q_j, m_j)$$

with

$$(64) \quad \begin{cases} \alpha_j(x) = \cos(b_j x) (z_1 \alpha_{j-1}(x) + z_2 \beta_{j-1}(x)) \\ \quad + \sin(b_j x) (-z_2 \alpha_{j-1}(x) - z_4 \beta_{j-1}(x)), \\ \beta_j(x) = \cos(b_j x) (z_2 \alpha_{j-1}(x) + z_4 \beta_{j-1}(x)) \\ \quad + \sin(b_j x) (z_1 \alpha_{j-1}(x) + z_2 \beta_{j-1}(x)). \end{cases}$$

That proves (59). Thanks to (64), we compute $W_j(x)$ and we find:

$$\begin{aligned}
W_j(x) &= b_j [(z_1^2 + z_2^2) \alpha_{j-1}^2(x) + 2z_2(z_1 + z_4) \alpha_{j-1}(x) \beta_{j-1}(x) + (z_2^2 + z_4^2) \beta_{j-1}^2(x)] \\
&+ (z_1 z_4 - z_2^2) W_{j-1}(x).
\end{aligned}$$

Since $(z_1^2 + z_2^2)(z_2^2 + z_4^2) - [z_2(z_1 + z_4)]^2 = (z_2^2 - z_1 z_4)^2 \geq 0,$ we deduce from the previous identity that

$$W_j(x) \geq (z_1 z_4 - z_2^2) W_{j-1}(x).$$

From (63) we find that

$$(z_1 z_4 - z_2^2) = \frac{m_j q_{j-1} (q_{j-1} + q_j)^2 (m_j q_{j-1}^2 + m_{j-1} q_j^2)^2}{m_{j-1}^3 q_j^7} > 0$$

Due to these last two inequalities, we get the conclusion :

$$d_j = (z_1 z_4 - z_2^2) d_{j-1} > 0. \quad \blacksquare$$

Lemma 4.9 (*asymptotic behaviour of the characteristic equation*)

Assume that the characteristic equation is still given by Theorem 4.5. Then there exists a constant K which is independent of the variable λ such that:

$$f(\sqrt{\lambda}) = e^{K\sqrt{\lambda}} \cdot (f_\infty(\sqrt{\lambda}) + g(\sqrt{\lambda}))$$

where

$$(65) \quad f_\infty(\sqrt{\lambda}) = e_4^t \cdot \left(\prod_{i=N}^1 H(q_i, b_i, m_i) \right) \cdot e_4$$

with $e_4^t = (0, 0, 0, 1, 0, 0)$, $H(q_i, b_i, m_i)$, $V_1(q, m)$ and $V_2(q, m)$ given in Definition 4.6. The function g satisfies $\lim_{\lambda \rightarrow +\infty} g(\sqrt{\lambda}) = 0$.

Thus, the asymptotic behaviour of the spectrum $\sigma(\mathcal{A})$ corresponds to the roots of the asymptotic characteristic equation

$$(66) \quad f_\infty(\sqrt{\lambda}) = 0.$$

These roots are all simple. Moreover, there exists a constant $d > 0$ (which depends only on the material constants) such that for any root x_0 of f_∞

$$(67) \quad |f'_\infty(x_0)| \geq d.$$

Proof.

- The first step is to prove the existence of the form $e^{K\sqrt{\lambda}} \cdot (f_\infty(\sqrt{\lambda}) + g(\sqrt{\lambda}))$ for $f(\sqrt{\lambda})$: each $\text{ext}(A_j)$ is of the form $\text{ext}(A_j) = e^{b_j \sqrt{\lambda}} H(q_j, b_j, m_j) + o(e^{b_j \sqrt{\lambda}})$ (cf. Lemma 4.7) with $H(q, b, m) = O(1)$.

Multiplying these expressions where j varies between 1 and N , we get:

$$e^{K\sqrt{\lambda}} \cdot \left(\prod_{j=N}^1 H(q_j, b_j, m_j) \right) + o(K\sqrt{\lambda}) \text{ with } K = \sum_{i=1}^N b_i.$$

- The second step is to compute $f_\infty(\sqrt{\lambda}) = e_4^t \cdot \left(\prod_{i=N}^1 H(q_i, b_i, m_i) \right) \cdot e_4$ using Lemma 4.8 and its proof. Since $e_4^t \cdot V_1(q, m) = 1$ and $e_4^t \cdot V_2(q, m) = 0$

$$(68) \quad f_\infty(\sqrt{\lambda}) = e_4^t \cdot H(q_N, b_N, m_N) \cdot \left(\prod_{i=N-1}^1 H(q_i, b_i, m_i) \right) \cdot e_4 = f_N c_N + g_N s_N$$

with $f_N = z_1 \alpha_{N-1} + z_2 \beta_{N-1}$ and $g_N = -z_2 \alpha_{N-1} - z_4 \beta_{N-1}$. And the wronskian $W(f_N, g_N)$ satisfies: $W(f_N, g_N) = (z_2^2 - z_1 z_4) W_{N-1} \leq d < 0$ with $d = (z_2^2 - z_1 z_4) d_{N-1}$ and d_{N-1} the constant introduced in Lemma 4.8 (depending only on the material constants).

- The third step is to compute the derivative of $f_\infty(x)$ (for the sake of completeness, we give the proof which is exactly the same one as in [16]).

$$f'_\infty(x) = \cos(b_N x)[f'_N(x) + b_N g_N(x)] + \sin(b_N x)[g'_N(x) - b_N f_N(x)].$$

We deduce that for all $x \in \mathbb{R}$, $\Delta(x) = (f_\infty(x))^2 + (f'_\infty(x))^2$ has the following form:

$$(69) \quad \Delta(x) = (\cos(b_N x) \sin(b_N x)) M(x) \begin{pmatrix} \cos(b_N x) \\ \sin(b_N x) \end{pmatrix},$$

where the matrix $M(x)$ is symmetric, positive and given by

$$M(x) = \begin{pmatrix} M_{11}(x) & M_{12}(x) \\ M_{21}(x) & M_{22}(x) \end{pmatrix}$$

and

$$\begin{cases} M_{11}(x) = f_N(x)^2 + b_N^2 g_N(x)^2 + 2b_N g_N(x) f'_N(x) + f'_N(x)^2, \\ M_{12}(x) = (1 - b_N^2) f_N(x) g_N(x) - b_N (f_N(x) f'_N(x) - g_N(x) g'_N(x)) + f'_N(x) g'_N(x), \\ M_{21}(x) = M_{12}(x), \\ M_{22}(x) = b_N^2 f_N(x)^2 + g_N(x)^2 - 2b_N f_N(x) g'_N(x) + g_N'(x)^2. \end{cases}$$

Let $\lambda_{min}(x), \lambda_{max}(x)$ be the two eigenvalues of $M(x)$ such that $0 \leq \lambda_{min}(x) \leq \lambda_{max}(x)$. After some computation we find

$$\begin{aligned} \lambda_{min}(x) \lambda_{max}(x) &= \det(M(x)) \\ &= b_N^2 (f_N(x)^2 + g_N(x)^2)^2 - 2b_N (f_N(x)^2 + g_N(x)^2) W(f_N, g_N)(x) \\ &\quad + W(f_N, g_N)(x)^2. \\ &= (W(f_N, g_N)(x) + b_N (f_N(x)^2 + g_N(x)^2))^2. \end{aligned}$$

Consequently with the estimate of the wronskian given at the end of the first step,

$$(70) \quad \forall x \in \mathbb{R}, \det(M(x)) = \lambda_{min}(x) \lambda_{max}(x) \geq W(f_N, g_N)(x)^2 \geq d^2.$$

On the other hand, since f_N and g_N are trigonometric polynomials, the trace of $M(x)$ is bounded on \mathbb{R} . Thus, there exists $d' > 0$ such that

$$(71) \quad \forall x \in \mathbb{R}, 0 \leq \text{tr}(M(x)) = \lambda_{min}(x) + \lambda_{max}(x) \leq d'^2.$$

From (70) and (71) we deduce that $\lambda_{min}(x) \geq \left(\frac{d}{d'}\right)^2 > 0$. Therefore from (69) we get

$$\forall x \in \mathbb{R}, \Delta(x) \geq \left(\frac{d}{d'}\right)^2 > 0.$$

That means that if x_0 is a root of f_∞ then $|f'_\infty(x_0)| \geq \frac{d}{d'} > 0$. ■

All the required properties are now proved to state the main result of this section:

Theorem 4.10 (*the spectral gap*)

Let λ_k^2 , $k \in \mathbb{N}^*$, ($\lambda_k > 0$) be the (strictly) monotone increasing sequence of eigenvalues of Problem (EP) given at the beginning of Section 4 then

$$(72) \quad \lim_{k \rightarrow +\infty} (\lambda_{k+1} - \lambda_k) = +\infty.$$

Proof. Since all the roots of f_∞ are simple and since there exists a constant $d > 0$ (which depends only on the material constants) such that for any root x_0 of f_∞

$$(73) \quad |f'_\infty(x_0)| \geq d.$$

it holds $\sqrt{\lambda_{k+1}} - \sqrt{\lambda_k} > \sigma$ with $\sigma > 0$ (cf. Theorem 5.3 of [14]).

Now $\lambda_{k+1} - \lambda_k = (\sqrt{\lambda_{k+1}} - \sqrt{\lambda_k})(\sqrt{\lambda_{k+1}} + \sqrt{\lambda_k})$ with $\lim_{k \rightarrow +\infty} \lambda_k = +\infty$, hence the announced result. ■

5 Proof of a uniform estimate for the eigenfunctions

Lemma 5.1 (*uniform estimate for $|\phi_N(l_N)|$*)

Consider the eigenvalue problem (EP) given in Section 4. For any eigenfunction ϕ associated to the eigenvalue λ^2 , there exists a constant K_1 such that:

$$(74) \quad K_1 \cdot \|\phi\|_{\mathcal{H}}^2 \leq |\phi_N(l_N)|^2$$

with the norm $\|\cdot\|_{\mathcal{H}}$ introduced in Section 2.3.1.

The proof of the lemma requires some technical intermediate results.

Let us first introduce some useful notation for the following.

Notation.

Keeping the notation a_j and l_j introduced in Section 2 as well as $q_j = (m_j/a_j)^{1/4}$, $b_j = q_j l_j$ introduced in Section 4.1, consider the functions $h_i(a_j, b_j, \lambda, x)$ for $i \in \{1; 2; 3; 4\}$ and $x \in [0; l_j]$ denoted $h_i(x)$ for the sake of simplicity:

$$(75) \quad \begin{cases} h_1(x) = \cos(q_j \sqrt{\lambda} x), \\ h_2(x) = \sin(q_j \sqrt{\lambda} x), \\ h_3(x) = \exp(-b_j \sqrt{\lambda}) \exp(q_j \sqrt{\lambda} x), \\ h_4(x) = \exp(-q_j \sqrt{\lambda} x). \end{cases}$$

$G(b_j, q_j)$ is the 4×4 Gram matrix defined by $(G(b_j, q_j))_{i,k} = \int_0^{l_j} h_i(x) h_k(x) dx$.

At last the matrices D and B_1 are:

$$(76) \quad D(a, q) = \frac{1}{4} \begin{pmatrix} 2 & 0 & -\frac{2}{aq^2} & 0 \\ 0 & \frac{2}{q} & 0 & -\frac{2}{aq^3} \\ 1 & -\frac{1}{q} & \frac{1}{aq^2} & -\frac{1}{aq^3} \\ e^{b\sqrt{\lambda}} & \frac{e^{b\sqrt{\lambda}}}{q} & \frac{e^{b\sqrt{\lambda}}}{aq^2} & \frac{e^{b\sqrt{\lambda}}}{aq^3} \end{pmatrix}$$

$$(77) \quad B_1(q, m) = \frac{1}{4} \begin{pmatrix} 1 & \frac{1}{q} & \frac{q^2}{m} & \frac{q}{m} \\ q & 1 & \frac{q^3}{m} & \frac{q^2}{m} \\ \frac{m}{q} & \frac{m}{q} & 1 & \frac{1}{q} \\ \frac{q^2}{m} & \frac{q^3}{m} & q & 1 \end{pmatrix}.$$

Lemma 5.2

Any eigenfunction ϕ associated to the eigenvalue λ^2 for the eigenvalue problem (EP) given at the beginning of Section 4 may be uniquely written as a linear combination of the (h_i) 's. Denote by $(C_j)_i$ the coefficients of the decomposition of ϕ_j in the basis $(h_i)_{i \in \{1;2;3;4\}}$

i.e. $\phi_j(x) = \sum_{i=1}^4 (C_j)_i h_i(x)$ for $j \in \{1, \dots, N\}$ and $x \in [0, l_j]$. Then

$$(78) \quad C_j = D(a_j, q_j)V_j(0), \text{ and } A(q, b, m) = \frac{1}{4} \exp(b\sqrt{\lambda})B_1(q, m)$$

with V_j and $A(q, b, m)$ defined in Section 4.1.

(V_j being computed for the j -th component of the particular eigenfunction ϕ).

There exists a positive constant C (by constant we mean independent of λ) such that

$$\|\phi\|_{\mathcal{H}}^2 = \sum_{j=1}^N \int_0^{l_j} |\phi_j(x)|^2 dx \leq C \max_{j \in \{1 \dots N\}} (C_j^t C_j).$$

Proof. Proving that the h_i 's are linearly independent is a classical computation. (78) is proved by calculation.

By definition of the inner product in \mathcal{H} (section 2.3.1):

$$\|\phi\|_{\mathcal{H}}^2 = \sum_{j=1}^N \int_0^{l_j} \phi_j(x)^2 dx = \sum_{j=1}^N C_j^t G(b_j, q_j) C_j.$$

Now, after calculation, the matrix $G(b, q)$ is:

$$(79) \quad G(b, q) = \frac{1}{2q\sqrt{\lambda}} \times \begin{pmatrix} cs + b\sqrt{\lambda} & s^2 & 1 - (c-s)e^{-b\sqrt{\lambda}} & c + s - e^{-b\sqrt{\lambda}} \\ s^2 & -cs + b\sqrt{\lambda} & 1 - (c+s)e^{-b\sqrt{\lambda}} & -c + s + e^{-b\sqrt{\lambda}} \\ 1 - (c-s)e^{-b\sqrt{\lambda}} & 1 - (c+s)e^{-b\sqrt{\lambda}} & 1 - e^{-2b\sqrt{\lambda}} & 2be^{-b\sqrt{\lambda}}\sqrt{\lambda} \\ c + s - e^{-b\sqrt{\lambda}} & -c + s + e^{-b\sqrt{\lambda}} & 2be^{-b\sqrt{\lambda}}\sqrt{\lambda} & 1 - e^{-2b\sqrt{\lambda}} \end{pmatrix}$$

with the notation $c = \cos(b\sqrt{\lambda})$, $s = \sin(b\sqrt{\lambda})$.

Note that all its terms are bounded with respect to λ . The estimate of $\|\phi\|_{\mathcal{H}}^2$ follows. ■

Lemma 5.3 (*estimate of $\phi_N(l_N)$*)

Let $M(\lambda)$ the 4×4 matrix defined by (48) in Section 4.1. Denote by $\alpha(\lambda)$ (respectively $\beta(\lambda)$) the first (resp. second) term of the first line of $M(\lambda)$ i.e.

$$\begin{cases} \alpha(\lambda) = e_1^t M(\lambda) e_1, \\ \beta(\lambda) = e_1^t M(\lambda) e_2, \end{cases}$$

with $e_1 = (1, 0, 0, 0)$ and $e_2 = (0, 1, 0, 0)$.

Then the eigenfunction ϕ of Problem (EP) associated to the eigenvalue λ^2 can be chosen such that $\phi_N(l_N) = \beta(\lambda)$ and the asymptotic behaviour of $\alpha(\lambda)$ and $\beta(\lambda)$ is given by:

$$\begin{cases} \alpha(\lambda) = C \cdot q_N \cdot e^{B\sqrt{\lambda}} + o\left(e^{B\sqrt{\lambda}}\right), \\ \beta(\lambda) = C \cdot e^{B\sqrt{\lambda}} + o\left(e^{B\sqrt{\lambda}}\right), \end{cases}$$

with $B := \sum_{j=1}^N b_j$.

Note that the constants C are not identical nor equal to those of Lemma 5.2 but we will always call the constants C . All of them are independent of λ but depend on the material constants given by the a_j 's, b_j 's...

Proof. Any eigenfunction ϕ associated to the eigenvalue λ^2 satisfies condition (19) of Section 2.3.1. In particular $\phi_1(0) = \phi_1'(0) = 0$ so the first two components of the vector $V_1(0)$ associated to ϕ (defined in Section 4.1) vanish. Moreover (25) and (26) also imply $\phi_N''(l_N) = 0$ and $\phi_N^{(3)}(l_N) = 0$ so the third and fourth components of $V_N(l_N)$ vanish : $V_N(l_N)$ is of the form $((V_N(l_N))_1, (V_N(l_N))_2, 0, 0)^t$.

Now, due to Lemma 4.1, $V_N(l_N) = M(\lambda)V_1(0)$ or $V_1(0) = M(\lambda)^{-1}V_N(l_N)$. Thus, if $\mu(\lambda)$ (respectively $\nu(\lambda)$) is the third (resp. fourth) term of the first line of $M(\lambda)^{-1}$, then $\mu(\lambda)(V_N(l_N))_1 + \nu(\lambda)(V_N(l_N))_2 = 0$.

$(V_N(l_N))_1 = \nu(\lambda)$ and $(V_N(l_N))_2 = -\mu(\lambda)$ is a solution of this equation which means that the eigenfunction ϕ of Problem (EP) associated to the eigenvalue λ^2 can be chosen such that $\phi_N(l_N) = \nu(\lambda)$ (such an eigenfunction is not normalized).

Now, to avoid the use of the inverse of the matrix $M(\lambda)$, we choose to switch the indices, which is equivalent to switch the boundary conditions i.e. in that proof as well as in the proofs of Lemmas 5.4 and 5.6, the definitions of V and $D(A)$ change : conditions (19), (25) and (26) of Section 2.3.1 become $\phi_N(l_N) = \phi_N'(l_N) = 0$ and $\phi_1''(0) = \phi_1^{(3)}(0) = 0$. Thus the eigenfunction ϕ of Problem (EP) associated to the eigenvalue λ^2 can be chosen such that $\phi_N(l_N) = \beta(\lambda)$.

The second part of the proof contains the estimate of some terms of the matrix $M(\lambda)$. Recall that

$$M(\lambda) = A_N A_{N-1} \dots A_2 A_1.$$

The asymptotic behaviour of $A_j = A(q_j, b_j, m_j)$ is given by $A_j = \exp(b_j \sqrt{\lambda}) B_1(q_j, m_j) + o(\exp(b_j \sqrt{\lambda}))$ with $B_1(q_j, m_j)$ defined by the decomposition

$$A(q, b, m) = \sum_{\epsilon \in \{-1; 0; 1\}} \exp(\epsilon b \sqrt{\lambda}) B_\epsilon(q, m).$$

Combining the above estimates and using (48) lead to the desired result. ■

The aim is still the estimation of $\|\phi\|_{\mathcal{H}}$ which requires, due to Lemma 5.2, the estimation of C_j .

Lemma 5.4 (*estimate of C_j*)

Let C_j be the vector already defined by (78) with a fixed $j \in \{1, \dots, N\}$, the vector \vec{b} be defined by $\vec{b} = (b_1, \dots, b_N)$ and denote by $\vec{u} \cdot \vec{v} := \sum_{i=1}^N u_i v_i$, then there exist vectors $W_{\vec{\epsilon}}(\lambda)$ such that C_j is of the form

$$(80) \quad C_j := \sum_{\vec{\epsilon} \in \{-2; -1; 0; 1\}^N} e^{\vec{b} \cdot \vec{\epsilon} \sqrt{\lambda}} W_{\vec{\epsilon}}(\lambda)$$

and all the terms of $W_{\vec{\epsilon}}(\lambda)$ are dominated by the function $\lambda \mapsto e^{B\sqrt{\lambda}}$ asymptotically (with $B := \sum_{j=1}^N b_j$).

More precisely the terms of $W_{\vec{\epsilon}}(\lambda)$ only contain expressions of the form $\cos(b_j \sqrt{\lambda})$ and $\sin(b_j \sqrt{\lambda})$ with $j \in \{1, \dots, N\}$.

Proof.

First Part. For a fixed $j \in \{1, \dots, N\}$, we start with isolating the terms containing $e^{b_j \sqrt{\lambda}}$ in the involved matrices.

$$\begin{cases} D(a_j, q_j) = e^{b_j \sqrt{\lambda}} D^+(a_j, q_j) + D^r(a_j, q_j), \\ A_j = A(q_j, b_j, m_j) = e^{b_j \sqrt{\lambda}} B^+(q_j, m_j) + B^r(q_j, m_j). \end{cases}$$

The decomposition of A_j is the same one as in the proof of Lemma 5.3 i.e. the matrix called B^+ in that proof is B_1 . The exponent r is chosen for the rest which does not contain $e^{b_j \sqrt{\lambda}}$.

Since $M(\lambda) = A_N \dots A_j \dots A_2 A_1$ (cf. (48)), it holds:

$$\begin{aligned} M(\lambda) &= A_N \dots A_{j+1} \left(e^{b_j \sqrt{\lambda}} B^+(q_j, m_j) + B^r(q_j, m_j) \right) A_{j-1} \dots A_2 A_1 \\ &= e^{b_j \sqrt{\lambda}} A_N \dots A_{j+1} B^+(q_j, m_j) A_{j-1} \dots A_2 A_1 + A_N \dots A_{j+1} B^r(q_j, m_j) A_{j-1} \dots A_2 A_1 \\ &=: e^{b_j \sqrt{\lambda}} M^+(\lambda) + M^r(\lambda). \end{aligned}$$

The first and second terms of the first line of the matrix $M(\lambda)$ denoted by α and β in Lemma 5.3 can be decomposed as follows:

$$\begin{cases} \alpha(\lambda) = e_1^t M(\lambda) e_1 = e^{b_j \sqrt{\lambda}} e_1^t M^+(\lambda) e_1 + e_1^t M^r(\lambda) e_1 = e^{b_j \sqrt{\lambda}} \alpha^+(\lambda) + \alpha^r(\lambda), \\ \beta(\lambda) = e_1^t M(\lambda) e_2 = e^{b_j \sqrt{\lambda}} \beta^+(\lambda) + \beta^r(\lambda). \end{cases}$$

Thus the vector $V_1(0)$ is decomposed as well: $V_1(0) = (\beta(\lambda), -\alpha(\lambda), 0, 0)^t = e^{b_1 \sqrt{\lambda}} V_1^+(0) + V_1^r(0)$ with $V_1^+(0) = (\beta^+(\lambda), -\alpha^+(\lambda), 0, 0)^t$. Then $C_j = D(a_j, q_j) A_{j-1} \cdots A_1 V_1(0)$ may be written as:

$$(81) \quad C_j = e^{2b_j \sqrt{\lambda}} C_j^{++} + e^{b_j \sqrt{\lambda}} C_j^+ + C_j^r \text{ with } C_j^{++} := D^+(a_j) A_{j-1} \cdots A_1 V_1^+(0)$$

where neither C_j^{++} , nor C_j^+ , nor C_j^r contains $e^{b_j \sqrt{\lambda}}$. The vanishing of C_j^{++} remains to be proved in order to establish (80).

Second Part. For a fixed $j \in \{1, \dots, N\}$, let us prove that $C_j^{++} = 0$ with C_j^{++} defined by (81). Recall that $M(\lambda) = e^{b_j \sqrt{\lambda}} M^+(\lambda) + M^r(\lambda)$ with $M^+(\lambda) = A_N \cdots A_{j+1} B^+(q_j, m_j) A_{j-1} \cdots A_2 A_1$. The matrices A_i for $i \in \{1, \dots, j-1\}$ defined in Section 4.1 are all invertible since their determinant is equal to 1 (calculation).

The matrix $B^+(q_j, m_j)$ is defined as follows:

$$(82) \quad B^+(q, m) = \frac{1}{4} \begin{pmatrix} 1 & \frac{1}{q} & \frac{q^2}{m} & \frac{q}{m} \\ q & 1 & \frac{q^3}{m} & \frac{q^2}{m} \\ \frac{m}{q} & \frac{m}{q} & 1 & \frac{1}{q} \\ \frac{q^2}{m} & \frac{q^3}{m} & q & 1 \end{pmatrix}.$$

Note that the columns of $B^+(q_j, m_j)$ are all proportional to the first one so the rank of $B^+(q_j, m_j)$ is 1. Thus the rank of $M^+(\lambda)$ is also 1 which means in particular that all its lines are proportional to the first one.

Now the first (respectively second) term of the first line of $M^+(\lambda)$ is, by definition, $\alpha^+(\lambda)$ (resp. $\beta^+(\lambda)$) and $V_1^+(0) = (\beta^+(\lambda), -\alpha^+(\lambda), 0, 0)^t$. So the first term of the product $M^+(\lambda) V_1^+(0)$ is 0. And since the other lines of $M^+(\lambda)$ are proportional to the first one, the other terms also vanish i.e. $M^+(\lambda) V_1^+(0) = 0$.

It is equivalent to $A_N \cdots A_{j+1} B^+(q_j, m_j) A_{j-1} \cdots A_2 A_1 V_1^+(0) = 0$ and, since $(A_N \cdots A_{j+1})$ is invertible, it implies: $A_{j-1} \cdots A_2 A_1 V_1^+(0) \in \text{Ker}(B^+(q_j, m_j))$.

The matrix $D^+(a_j, q_j)$ is defined as follows:

$$D^+(a_j, q_j) = \frac{1}{4} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & \frac{1}{q_j} & \frac{1}{a_j q_j^2} & \frac{1}{a_j q_j^3} \end{pmatrix}.$$

Since $q_j^2/m_j = 1/(a_j q_j^2)$, it clearly holds $\text{Ker}(B^+(q_j, m_j)) = \text{Ker}(D^+(a_j, q_j))$.

Thus $C_j^{++} := D^+(a_j, q_j) A_{j-1} \cdots A_2 A_1 V_1^+(0) = 0$. ■

Lemma 5.5 (estimate of $V_j(l_j)$)

Let $j \in \{1, \dots, N\}$ and let the vector V_j be defined as in Section 4.1. For any $K > 0$, there exists a positive constant C (independent of λ) such that, if $\lambda > K$, for $j \in \{1, \dots, N\}$ and $i \in \{1; 2; 3; 4\}$

$$(83) \quad |e_i^t V_j(l_j)| \leq C e^{B\sqrt{\lambda}}$$

with $e_1^t = (1, 0, 0, 0)$, $e_2^t = (0, 1, 0, 0)$, $e_3^t = (0, 0, 1, 0)$, $e_4^t = (0, 0, 0, 1)$ and $B := \sum_{j=1}^N b_j$.

The constants are still all denoted by C . They only depend on the material constants.

Proof. The property is proved by induction. The vector $V_1(0)$ is: $V_1(0) = (\beta(\lambda), -\alpha(\lambda), 0, 0)^t$ and the behaviour of α and β is given by Lemma 5.3. The components of $V_1(l_1) = A_1 V_1(0)$ keep the same fastest growing term as the terms of $V_1(0)$. Indeed the multiplication by the matrix A_1 which contains exponential terms could a priori change the exponential into $e^{(2b_1 + \sum_{j=2}^N b_j)\sqrt{\lambda}}$ but as it was proved for C_j in the proof of Lemma 5.4, it is not the case. ■

Lemma 5.6 (a more precise estimate of C_j)

Let C_j be the vector already defined by (78) with a fixed $j \in \{1, \dots, N\}$. For any $K > 0$, there exists a constant C (independent of λ) such that, if $\lambda > K$ and $i \in \{1; 2; 3; 4\}$:

$$(84) \quad |(C_j)_i| \leq C e^{B\sqrt{\lambda}}$$

with $(C_j)_i$ the i -th term of the vector C_j as in Lemma 5.2 and $B := \sum_{j=1}^N b_j$.

Proof. Recall that $C_j = D(a_j, q_j) V_j(0)$ (cf. (78)). We have just proved in the second part of the proof of Lemma 5.5 that, for any $j \in \{1, \dots, N\}$, the absolute values of the four terms of $V_{j+1}(0)$ are bounded by $C e^{B\sqrt{\lambda}}$ for large values of λ . It is also clear for $V_1(0) = (\beta(\lambda), -\alpha(\lambda), 0, 0)^t$ due to the estimates of α and β given in Lemma 5.3.

Now the matrix $D(a_j, q_j)$ contains exponential terms but we proved in the proof of Lemma 5.4 that they do not affect the fastest growing term of C_j . Hence the result. ■

Theorem 5.7 (first uniform estimate for $|\phi_N(l_N)|$)

Consider the eigenvalue problem (EP) given in Section 3.1. For any eigenfunction $\phi \in D(A)$ associated to the eigenvalue λ^2 , there exist a constant K_1 such that:

$$(85) \quad K_1 \cdot \|\phi\|_{\mathcal{H}}^2 \leq |\phi_N(l_N)|^2$$

with the norm $\|\cdot\|_{\mathcal{H}}$ introduced in Section 2.3.1.

Proof. Due to Lemma 5.2

$$\|\phi\|_{\mathcal{H}}^2 \leq C \max_{j \in \{1 \dots N\}} (C_j^t C_j).$$

Then (84) implies $\|\phi\|_{\mathcal{H}}^2 \leq C \left(e^{B\sqrt{\lambda}} \right)^2$.

Now, we stated in the proof of Theorem 5.3 that the eigenfunction ϕ of Problem (EP) associated to the eigenvalue λ^2 can be chosen such that $\phi_N(l_N) = \beta(\lambda)$. The estimate of $\beta(\lambda)$ for large values of λ given by Theorem 5.3 gives the desired result. ■

5.1 Second estimate: admissibility

Theorem 5.8 (*second estimate for controllability*)

Consider the eigenvalue problem (EP) associated to Problem (P) (given in Section 4). For any eigenfunction ϕ associated to the eigenvalue λ^2 , there exists a constant K_2 such that:

$$(86) \quad |\phi_N(l_N)|^2 \leq K_2 \cdot \|\phi\|_{\mathcal{H}}^2$$

with the norm $\|\cdot\|_{\mathcal{H}}$ defined in Section (2.3.1).

Proof. We established in the proof of Lemma 5.2

$$\|\phi\|_{\mathcal{H}}^2 = \sum_{j=1}^N \int_0^{l_j} \phi_j(x)^2 dx = \sum_{j=1}^N C_j^t G(b_j, q_j) C_j$$

with C_j and $G(b_j, q_j)$ defined in the same Lemma. Thus $\|\phi\|_{\mathcal{H}}^2 \geq C_1^t G(b_1, q_1) C_1$ and it remains to estimate this expression from below.

Due to (78) it holds $C_1 = D(a_1, q_1)V_1(0)$ and we stated in Lemma 5.3 that

$$V_1(0) = (\beta(\lambda), -\alpha(\lambda), 0, 0)^t = (\beta, q_N \cdot \beta + o(\beta), 0, 0)^t$$

as λ and thus β tend to infinity. Then, multiplying by the matrix $D(a_1, q_1)$ given just before Lemma 5.2, it follows

$$C_1 = 2 \left(\beta, \beta + \frac{o(\beta)}{q_1}, -\frac{o(\beta)}{2q_1}, \beta e^{b_1 \sqrt{\lambda}} + \frac{e^{b_1 \sqrt{\lambda}}}{2q_1} \cdot o(\beta) \right).$$

Now we proved in Lemma 5.6 that, for any $K > 0$, there exists a constant C (independent of λ) such that, if $\lambda > K$ and $i \in \{1; 2; 3; 4\}$, then $|(C_j)_i| \leq C\beta(\lambda)$ with $(C_j)_i$ the i -th term of the vector C_j . So the fourth term of C_1 grows as fast as β i.e.

$$C_1 = 2 \left(\beta, \beta + \frac{o(\beta)}{q_1}, -\frac{o(\beta)}{2q_1}, O(\beta) \right).$$

Looking thoroughly at the terms of the matrix $G(b_1, q_1)$ given in the proof of Lemma 5.2, we can see that only two terms do not tend to zero as λ tends to infinity which can be written as:

$$G(b_1, q_1) = \begin{pmatrix} \frac{b_1}{2q_1} & 0 & 0 & 0 \\ 0 & \frac{b_1}{2q_1} & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} + o(1).$$

It follows $C_1^t \cdot G(b_1, q_1) \cdot C_1 = \frac{4b_1}{q_1} \cdot \beta^2 + o(\beta^2)$ and since $|\phi_N(l_N)|^2 = |\beta(\lambda)|^2$, the desired estimate follows. ■

Acknowledgement.

The authors thank the referees for their attentive reading of the manuscript and the questions they asked.

References

- [1] F. Ali Mehmeti. *Nonlinear Waves in Networks*. Akademie Verlag, Berlin, 1994.
- [2] F. Ali Mehmeti. *Transient Waves in Semi-Infinite Structures : the Tunnel Effect and the Sommerfeld Problem*. Akademie Verlag, Berlin, 1996.
- [3] F. Ali Mehmeti, J. von Below and S. Nicaise, edit. *Partial differential equations on multi-structures, Lecture Notes in Pure and Applied Mathematics*, Marcel Dekker, 2001.
- [4] K. Ammari, M. Tucsnak. *Stabilization of second order evolution equations by a class of unbounded feedbacks*. ESAIM Control Optim. Calc. Var. **6**, p. 361-386, 2001.
- [5] K. Ammari, M. Mehrenberger. *Stabilization of coupled systems*. Acta Math. Hungar. **123** (1-2), p. 1-10, 2009.
- [6] K. Ammari, D. Mercier, V. Régnier, J. Valein. *Spectral analysis and stabilization of a chain of serially connected Euler-Bernoulli beams and strings*. Comm. In Pure and Appl. Anal. **11/2**, p. 785-807, 2012.
- [7] W. Arendt and C. J. K. Batty. *Tauberian theorems and stability of one-parameter semi-groups*. Trans. Amer. Math. Soc., **305**, p. 837-852, 1988.
- [8] J. von Below. *A characteristic equation associated to an eigenvalue problem on C^2 -networks*. Linear Algebra Appl., **71**, p. 309-325, 1985.
- [9] C. Castro, E. Zuazua. *Analyse spectrale et contrôle d'un système hybride composé de deux poutres connectées par une masse ponctuelle*. C. R. Acad. Sci. Paris, **322**, p. 351-356, 1996.
- [10] G. Chen, M. Delfour, A. Krall, G. Payre. *Modeling, stabilization and control of serially connected beams*. SIAM J. Control Opt., **25**, p. 526-546, 1987.
- [11] B. Dekoninck, S. Nicaise. *The eigenvalue problem for network of beams*. Linear Algebra and its Applications, **314**, p. 165-189, 2000.
- [12] A. Haraux. *Séries lacunaires et contrôle semi-interne des vibrations d'une plaque rectangulaire*. J. Maths Pures et Appl., **68**, p. 457-465, 1989.
- [13] J.E. Lagnese, G. Leugering, E.J.P.G. Schmidt. *Modeling, Analysis and Control of Dynamic Elastic Multi-Link Structures*. Birkhäuser, Boston, 1994.
- [14] D. Mercier. *Spectrum analysis of a serially connected Euler-Bernoulli beams problem*. Netw. Heterog Media, **4**, p. 874-894, 2009.

- [15] D. Mercier, V. Régnier. *Spectrum of a network of Euler-Bernoulli beams*. J. Math. Anal. and Appl., **337/1**, p.174-196, 2007.
- [16] D. Mercier, V. Régnier. *Boundary controllability of a chain of serially connected Euler-Bernoulli beams with interior masses*. Collect. Math., **60/3**, p. 307-334, 2009.
- [17] S. Nicaise. *Diffusion sur les espaces ramifiés*. Thèse de doctorat, Mons, Belgique, 1986.
- [18] W. H. Paulsen. *The exterior matrix method for sequentially coupled fourth-order equations*. J. of Sound And Vibration 308, **4**, p. 132-163, 2007.