

The heat equation under conditions on the moments in higher dimensions

Delio Mugnolo, Serge Nicaise

▶ To cite this version:

Delio Mugnolo, Serge Nicaise. The heat equation under conditions on the moments in higher dimensions. Mathematical News / Mathematische Nachrichten, 2015, 288 (2-3), pp.295-308. $10.1002/\mathrm{mana}.201300298$. hal-03163722

HAL Id: hal-03163722 https://uphf.hal.science/hal-03163722

Submitted on 5 Jul2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE HEAT EQUATION UNDER CONDITIONS ON THE MOMENTS IN HIGHER DIMENSIONS

DELIO MUGNOLO AND SERGE NICAISE

ABSTRACT. We consider the heat equation on the N-dimensional cube $(0,1)^N$ and impose different classes of integral conditions, instead of usual boundary ones. Well-posedness results for the heat equation under the condition that the moments of order 0 and 1 are conserved had been known so far only in the case of N = 1 – for which such conditions can be easily interpreted as conservation of mass and barycenter. In this paper we show that in the case of general N the heat equation with such integral conditions is still well-posed, upon suitably relax the notion of solution. Existence and uniqueness of solutions with general initial data in a suitable space of distibutions over $(0, 1)^N$ are proved by introducing two appropriate realizations of the Laplacian and checking by form methods that they generate analytic semigroups. The solution thus obtained does however solve the heat equation only in a certain distributional sense. However, it turns out that one of these realizations is tightly related to a wellknown object of operator theory, the Krein-von Neumann extension of the Laplacian. This connection also establishes well-posedness in a classical sense, as long as the initial data are L^2 -functions.

1. INTRODUCTION

Fifty years ago, J.R. Cannon has suggested in [6] that realistic mechanical considerations suggest to study diffusion equations imposing a condition on the moment of order 0 of the unknown (i.e., on the total mass of the system), thus dropping one of the boundary conditions. Cannon's analysis was limited to the one-dimensional case, both in [6] and his later research surveyed in [7]. His original setting has been significantly generalized over the years (cf. [16, § 1] for a historical overview), in particular replacing also the remaining boundary condition by a condition on the moment of order 1 (i.e., on the barycenter).

Still, to the best of our knowledge only a few tentative extensions of the above conditions for heat or wave equations on higher dimensional domains have been proposed in the literature: We mention [15, 14, 17], where earlier related references are also collected. A first difficulty is that in case of a bounded domain in dimension $N \ge 2$ (unlike in the 1-dimensional case) infinitely many (boundary) conditions are necessary to determine a solution, while prescribing the moments of order 0 and 1 yields only finitely many conditions.

In this note, we introduce a general setting which in our opinion yields the proper N-dimensional extensions of the 1-dimensional moment conditions studied in [5, 16]. Because the moments of order 0 and 1 of a given function are its integrals against the two vectors that span the space of one-dimensional harmonic functions,

²⁰¹⁰ Mathematics Subject Classification. 47D06 (primary), 35J20 (secondary).

Key words and phrases. Non-local conditions; Analytic semigroups; Quadratic forms; Kreinvon Neumann extension.

This article has been initiated during a visit of the first author in Valenciennes and continued during a visit of the second author in Ulm. The authors thank the Université de Valenciennes et du Hainaut Cambrésis and the Universität Ulm for their hospitality. Both visits were financially supported by the Land Baden–Württemberg in the framework of the *Juniorprofessorenprogramm* – research project on "Symmetry methods in quantum graphs". We thank Fritz Gesztesy and Konstantin Pankrashkin for interesting discussions on the topic of Section 6.

it is natural to conjecture that for $N \geq 2$ Cannon's setting should be extended by imposing orthogonality to the harmonic functions. This allows to define two diffusion-type operators A and \tilde{A} and to study the associated parabolic problem.

We stress that neither of these operators is a classical Laplacian: Rather, each of them is an isomorphic image of (a realization of) the Laplacian in a suitable space of distributions of H^{-1} -type. An analogous problem has been already observed in [16]. Roughly speaking, A turns out to be an operator with conditions on the moment of order 0 along with uncountably many further conditions (that however impose in particular that the N linear moments of order 1 of the unknown have to vanish); whereas \tilde{A} will be an operator with periodic boundary conditions and a condition on the moment of order 0 only. This is discussed in detail in Section 4. Nevertheless, A and \tilde{A} agree with the Laplacian in the sense of distributions.

In the special case of N = 1 it has been shown in [16, §§ 3–4] that the solution of the abstract Cauchy problems associated with either of them also solves the classical heat equation. Indeed, the orbits of the semigroups generated by A and \tilde{A} contain only elements whose smoothness suffices to ensure that the generator acts on them as the usual second derivative: This can be proved carefully describing the smoothing enhancement yielded by analyticity of the semigroups.

It seems that a similar strategy is not successful in the present higher dimensional context. In fact, a direct computation suggests that in general the elements of the orbits are not much better than L^2 . Nevertheless, we will eventually show in Section 5 that the semigroups actually solve the usual heat equation (with moment conditions) – at least in a suitably weak, distributional sense.

Finally, in Section 6 we comment on the well-posedness of the relevant diffusion problem in a more usual L^2 -setting. We also emphasize the connections between our investigations and Krein's theory of self-adjoint extensions: Also this observation seems to be new.

2. The functional setting

Let $\Omega \subset \mathbb{R}^N$ be an open domain with Lipschitz boundary. It is well-known (see e.g. [1, Thm. 2.5]) that

(2.1)
$$H^{1}(\Omega) = H^{1}_{0}(\Omega) \oplus \operatorname{Har}(\Omega),$$

where $\operatorname{Har}(\Omega)$ denotes the space of (weakly) harmonic functions, i.e.,

$$\operatorname{Har}(\Omega) := \left\{ u \in H^1(\Omega) : \int_{\Omega} \nabla u \overline{\nabla v} dx = 0 \,\,\forall v \in H^1_0(\Omega) \right\}.$$

Furthermore, the trace operator γ_0 is an isomorphism between $\operatorname{Har}(\Omega)$ (the orthogonal of its null space) and $H^{\frac{1}{2}}(\partial\Omega)$ (its range).

If T^N denotes the N-dimensional torus, we can consider the (Hilbert) space $H^1(T^N)$, defined as usual as the space of periodic functions defined on T^N . Throughout this article we will denote

$$\mu_0(h) := \langle h, 1 \rangle, \qquad h \in H^{-1}(T^N),$$

the mean – i.e., the moment of order 0 – of a distribution h.

Now, $H_0^1((0,1)^N)$ is a closed subspace of $H^1(T^N)$, hence we can consider its orthogonal complement which, as above, turns out to be (2.2)

$$\operatorname{Har}(T^N) := \left\{ u \in H^1(T^N) : \int_{T^N} \nabla u \overline{\nabla v} dx = 0 \ \forall v \in H^1_0(\Omega) \right\} =: H^1(T^N) \ominus H^1_0\left((0,1)^N\right)$$

Hence, using $L^2((0,1)^N)$ as pivot space we can find the decomposition

(2.3)
$$H^{-1}(T^N) = H^{-1}\left((0,1)^N\right) \oplus \left(\operatorname{Har}(T^N)\right)',$$

where $H^{-1}((0,1)^N) := (H^1_0(0,1)^N)'.$

Remark 2.1. In other words, $\operatorname{Har}(T^N)$ is the space of periodic H^1 -functions that are (weakly) harmonic, whereas $\operatorname{Har}((0,1)^N)$ is the space of all (weakly) harmonic H^1 -function on the unit cube. It is by definition clear that $\operatorname{Har}(T^N)$ is a subspace of $\operatorname{Har}((0,1)^N)$. In fact, it will in general be a proper subspace: For N = 1 we e.g. find that $\operatorname{Har}(T^N)$ and $\operatorname{Har}((0,1)^N)$ are isomorphic to the spaces \mathbb{P}_0 and \mathbb{P}_1 of polynomials of degree 0 and of degree less than or equal to 1, respectively.

Let us denote by Id the orthogonal projector of $H^{-1}(T^N)$ onto its closed subspace $H^{-1}((0,1)^N)$ and by Id_m the restriction of Id to the annihilator

$$H := \{ w \in H^{-1}(T^N) : \langle w, v \rangle = 0 \ \forall v \in \operatorname{Har}(T^N) \}$$

of Har (T^N) , which is an isomorphism from H to $H^{-1}((0,1)^N)$.

Denote by

$$H^{\frac{1}{2}}(\partial T^{N}) := \{\gamma_{0}v : v \in H^{1}(T^{N})\},\$$

where γ_0 is here the trace operator from $H^1((0,1)^N)$ into $H^{\frac{1}{2}}(\partial(0,1)^N)$. The space $H^{\frac{1}{2}}(\partial T^N)$ is smaller than $H^{\frac{1}{2}}(\partial(0,1)^N)$ due to the periodicity assumption on elements from $H^1(T^N)$. Nevertheless $H^{\frac{1}{2}}(\partial T^N)$ is a Hilbert space with the induced norm

$$\|\varphi\|_{H^{\frac{1}{2}}(\partial T^{N})} := \inf_{\substack{v \in H^{1}(T^{N}) \\ \varphi = \gamma_{0}v}} \|v\|_{H^{1}(T^{N})}.$$

Lemma 2.2. For all $\varphi \in H^{\frac{1}{2}}(\partial T^N)$, there exists a unique $R\varphi \in Har(T^N)$ such that

 $\gamma_0 R \varphi = \varphi \ on \ \partial(0,1)^N.$

In other words, $\operatorname{Har}(T^N)$ is isomorphic to $H^{\frac{1}{2}}(\partial T^N)$.

The existence of the right inverse of the trace operator, seen as an operator from the space of weakly harmonic functions to the space of $H^{\frac{1}{2}}$ -functions over the boundary of a domain, is classical - see e.g. the much more general discussion in [13, § 2.7] or the alternative approach in [10, Lemma 1.2].

Define

$$\tilde{V} := \{ f \in L^2 \left((0,1)^N \right) : (f|g) = 0 \ \forall g \in \operatorname{Har}(T^N) \}, \\ V := \{ f \in L^2 \left((0,1)^N \right) : (f|g) = 0 \ \forall g \in \operatorname{Har} \left((0,1)^N \right) \},$$

that are clearly two closed subspaces of $L^2((0,1)^N)$. Indeed by introducing V_0 and V_1 as the closure of $\operatorname{Har}(T^N)$ and $\operatorname{Har}((0,1)^N)$ in $L^2((0,1)^N)$, respectively,

$$V_0 := \overline{\operatorname{Har}(T^N)}^{L^2}$$
$$V_1 := \overline{\operatorname{Har}((0,1)^N)}^{L^2}$$

one sees that \tilde{V} and V are the L^2 -orthogonal complements of V_0 and V_1 , respectively – shortly:

$$\tilde{V} = V_0^{\perp}$$
 and $V = V_1^{\perp}$

As $\operatorname{Har}(T^N) \subset \operatorname{Har}((0,1)^N)$ and hence $V_0 \subset V_1$, we conclude that

$$V \subset \tilde{V}.$$

We are going to need below a certain characterization of V_0 . For that purpose, we can notice that V_0 is trivially included into the domain of Δ_{L^2} – the part of Δ in $L^2((0,1)^N)$ –, i.e., into

$$D(\Delta_{L^2}) := \left\{ v \in L^2\left((0,1)^N \right) : \Delta v \in L^2\left((0,1)^N \right) \right\},\$$

which is a Hilbert space in its own right whenever equipped with the natural graph norm

$$v \mapsto \left(\|v\|_{L^2((0,1)^N)}^2 + \|\Delta v\|_{L^2((0,1)^N)}^2 \right)^{\frac{1}{2}}.$$

It follows from the general theory of interior elliptic regularity that if $u \in D(\Delta_{L^2})$, then in general one only has $u \in H^2_{\text{loc}}$, hence a trace of u need not exist as an element of $L^2(\partial(0,1)^N)$. However, we can give a meaning of its trace on each face as a vector in a suitable space of distributions. Indeed following [11, Thm 1.5.3.4], the space $\mathcal{D}([0,1]^N)$ – the set of the restriction of elements of $\mathcal{D}(\mathbb{R}^N)$ to $(0,1)^N$ – is dense in $D(\Delta_{L^2})$ and for all $i = 1, \dots, n$, the trace operator

$$\gamma_{i\pm}: v \mapsto v_{\Gamma_{i\pm}}$$

which is certainly defined for $v \in \mathcal{D}([0,1]^N)$, has a unique continuous extension from $D(\Delta_{L^2})$ into $(\tilde{H}^{\frac{1}{2}}(\Gamma_{i\pm}))'$. We denote also this extension by $\gamma_{i\pm}$. Here and below we are denoting by $\Gamma_{i\pm}$ the faces of the hypercube, which are defined by

$$\Gamma_{i-} := \{ x \in [0,1]^N : x_i = 0 \text{ and } x_j \in (0,1), \forall j \neq i \}, \\ \Gamma_{i+} := \{ x \in [0,1]^N : x_i = 1 \text{ and } x_j \in (0,1), \forall j \neq i \}.$$

Furthermore $\tilde{H}^{\frac{1}{2}}(\Gamma_{i\pm})$ denotes the subspace of elements $w \in H^{\frac{1}{2}}(\Gamma_{i\pm})$ such that \tilde{w} , its extension by zero outside $\Gamma_{i\pm}$, belongs to $H^{\frac{1}{2}}(\partial(0,1)^N)$.

Definition 2.3. Let $s > \frac{1}{2}$. We call a function $v \in D(\Delta_{L^2}) \cup H^s((0,1)^N)$ periodic if it satisfies

(2.4)
$$\gamma_{i-}v = T_i\gamma_{i+}v \text{ in } \left(\tilde{H}^{\frac{1}{2}}(\Gamma_{i+})\right)', \quad \forall i=1,\cdots,n.$$

Here $T_i: \left(\tilde{H}^{\frac{1}{2}}(\Gamma_{i+})\right)' \to \left(\tilde{H}^{\frac{1}{2}}(\Gamma_{i-})\right)'$ is the operator defined by means of

$$\langle T_i\psi,\varphi\rangle := \langle \psi,T_i^*\varphi\rangle \ \forall\psi\in \left(\tilde{H}^{\frac{1}{2}}(\Gamma_{i+})\right)',\varphi\in \tilde{H}^{\frac{1}{2}}(\Gamma_{i-}),$$

with

$$(T_i^*\varphi)(x_1,\cdots,1,\cdots,x_n) := \varphi(x_1,\cdots,0,\cdots,x_n)$$
 for a.e. $(x_1,\cdots,1,\cdots,x_n) \in \Gamma_{i+1}$
Lemma 2.4. One has

$$V_0 \subset \left\{ v \in L^2\left((0,1)^N\right) : \Delta v = 0 \text{ in } \mathcal{D}'\left((0,1)^N\right) \text{ satisfying } (2.4) \right\}.$$

Proof. As $\operatorname{Har}(T^N) \subset H^1(T^N)$, any $u \in \operatorname{Har}(T^N)$ clearly satisfies (2.4). Since there exists C > 0 such that

$$\|\gamma_{i-}v - T_{i}\gamma_{i+}v\| \le C \left(\|v\|_{L^{2}((0,1)^{N})}^{2} + \|\Delta v\|_{L^{2}((0,1)^{N})}^{2} \right)^{\frac{1}{2}}, \quad \forall v \in D(\Delta_{L^{2}}),$$

we directly conclude that any $v \in V_0$ still satisfies (2.4).

Remark 2.5. Note that in particular any function in $H^1(T^N)$ is periodic in the sense of the above definition, and by Lemma 2.4 so is any element of $H^1(T^N) + V_0$, too.

3. An equivalent inner product in $H^{-1}(T^N)$

Lemma 3.1. For all $f \in H^{-1}(T^N)$ there exists a unique $u_f \in H^1_m(T^N)$ such that $\operatorname{div} \nabla u_f = f - \mu_0(f).$

An equivalent inner product in $H^{-1}(T^N)$ is therefore given by

$$(\nabla u_f | \nabla u_g)_{L^2(T^N)^N} + \mu_0(f)\mu_0(\bar{g}), \qquad f, g \in H^{-1}(T^N).$$

In the proof we will need a closed subspace $H^1_m(T^N)$ of $H^1(T^N)$ defined by

$$H_m^1(T^N) := \left\{ u \in H^1(T^N) : \int_{(0,1)^N} u(x) dx = 0 \right\}$$

By the Poincaré-type inequality

$$\exists C > 0 : \|u\|_{L^{2}(T^{N})} \leq C \|\nabla u\|_{L^{2}(T^{N})^{N}}, \forall u \in H^{1}_{m}(T^{N}),$$

see e.g. [8, Prop. 5.44], which is based on the compactness of the embedding of $H^1((0,1)^N)$ into $L^2((0,1)^N)$, cf. [11, Thm. 1.4.3.2]), we know that

$$u \mapsto \|\nabla u\|_{L^2(T^N)^N}$$

defines a norm on $H^1_m(T^N)$ that is equivalent to the standard H^1 -norm.

Proof. Given $f \in H^{-1}(T^N)$, we set

$$f_0 = f - \langle f, 1 \rangle,$$

where here and below the duality bracket is between $H^{-1}(T^N)$ and $H^1(T^N)$. Hence $f_0 \in H^{-1}(T^N)$ and satisfies

$$\langle f_0, 1 \rangle = 0.$$

Now by the Theorem of Riesz–Fréchet there exists a unique solution $u_f \in H^1_m(T^N)$ of

(3.1)
$$\int_{(0,1)^N} \nabla u_f \cdot \nabla \bar{v} dx = -\langle f_0, v \rangle \ \forall v \in H^1_m(T^N).$$

Since $\langle f_0, 1 \rangle = 0$, this identity remains valid on the whole $H^1(T^N)$, namely

(3.2)
$$\int_{(0,1)^N} \nabla u_f \cdot \nabla \bar{v} dx = -\langle f_0, v \rangle \ \forall v \in H^1(T^N).$$

By choosing smooth enough test functions v, we see that

 $\operatorname{div} \nabla u_f = f_0 \text{ in } \mathcal{D}' \left((0, 1)^N \right),$

or equivalently

(3.3)
$$f = \operatorname{div} \nabla u_f + \langle f, 1 \rangle.$$

According to (3.2), we have

$$\|\nabla u_f\|_{L^2(T^N)^N}^2 = -\langle f_0, u_f \rangle \le \|f_0\|_{H^{-1}(T^N)} \|u_f\|_{H^1(T^N)},$$

and by the equivalence of norm mentioned before, we get

$$\|\nabla u_f\|_{L^2(T^N)^N} \lesssim \|f_0\|_{H^{-1}(T^N)} \lesssim \|f\|_{H^{-1}(T^N)}.$$

Conversely (3.2) is equivalent to

$$\langle f, v \rangle = -\int_{(0,1)^N} \nabla u_f \cdot \nabla \overline{v} dx + \langle f, 1 \rangle \overline{\langle v, 1 \rangle} \ \forall v \in H^1(T^N).$$

Consequently

$$\begin{split} \|f\|_{H^{-1}(T^{N})} &= \sup_{\|v\|_{H^{1}(T^{N})}=1} |\langle f, v \rangle| \\ &\leq \sup_{\|v\|_{H^{1}(T^{N})}=1} \left(\left| \int_{(0,1)^{N}} \nabla u_{f} \cdot \nabla \bar{v} dx \right| + |\langle f, 1 \rangle || \langle v, 1 \rangle| \right) \\ &\lesssim \|\nabla u_{f}\|_{L^{2}(T^{N})^{N}} + |\langle f, 1 \rangle|. \end{split}$$

This completely proves the assertion.

Remark 3.2. Let us briefly compare the theory we have just developed with that introduced in [16] in the one-dimensional setting. We define the "primitive" $P_N f \in L^2(T^N)^N$ of any $f \in H^{-1}(T^N)$ by

$$P_N f := \nabla u_f + \frac{\mu_0(f)}{N} \left(\vec{x} - \frac{\vec{1}}{2} \right), \quad \forall f \in H^{-1}(T^N)$$

where \vec{x} and $\frac{\vec{1}}{2}$ denote the vector-valued functions

$$\vec{x}: (x_1, \cdots, x_N) \mapsto \begin{pmatrix} x_1 \\ \vdots \\ x_N \end{pmatrix}$$
 and $\frac{\vec{1}}{2}: (x_1, \cdots, x_N) \mapsto \frac{1}{2} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}$,

respectively. From this expression, we see that

$$\operatorname{div} P_N f = f \text{ in } \mathcal{D}' \left((0,1)^N \right),$$

and

$$\langle P_N f, \alpha \rangle = 0 \ \forall \alpha \in \mathbb{C}^N.$$

It finally follows from Lemma 3.1 that the inner product

$$(f|g)_{H^{-1}(T^N)} = (P_N f|P_N g)_{L^2(T^N)} + \mu_0(f)\mu_0(\bar{g}),$$

induces a norm equivalent to the standard norm of $H^{-1}(T^N)$.

Note further that if $f \in L^2(T^N)$, then the solution $u_f \in H^1_m(T^N)$ of (3.1) belongs to $H^2(T^N)$ and therefore $P_N f$ belongs to $H^1(T^N)^N$.

Definition 3.3. We define an operator L from $H^1((0,1)^N)$ to H by

$$Lf := \mathrm{Id}_m^{-1}(\Delta f).$$

Observe that L is well-defined because $\Delta f \in H^{-1}((0,1)^N)$ for all $f \in H^1((0,1)^N)$.

Remark 3.4. If in particular $f \in H^1(T^N) \cap H^2((0,1)^N)$ satisfying Neumann boundary conditions, then $Lf = \Delta f \in L^2((0,1))^N \subset H^{-1}((0,1)^N)$ and hence in particular $\mathrm{Id}_m^{-1}(\Delta f) \in H_m^{-1}(T^N)$. Hence there exists by Lemma 3.1 a unique $u_{Lf} \in H_m^1(T^N)$ such that

$$\Delta u_{Lf} = Lf - \mu_0(Lf),$$

that is,

$$\Delta u_{Lf} = Lf.$$

Because $f - \mu_0(f) \in H^1_m(T^N)$, one can easily conjecture that

(3.4)
$$u_{Lf} = f - \mu_0(f)$$

This is indeed the case, as

$$\int_{(0,1)^N} \nabla(f - \mu_0(f)) \cdot \nabla \bar{v} \, dx = \int_{(0,1)^N} \nabla f \cdot \nabla \bar{v} \, dx$$
$$= \int_{\partial(0,1)^N} \frac{\partial f}{\partial n} \bar{v} dx - \int_{(0,1)^N} \Delta f \bar{v} \, dx$$
$$= -\int_{(0,1)^N} \Delta f \bar{v} \, dx, \quad \forall v \in H^1_m(T^N),$$

by the Gauß–Green formula.

Remark 3.5. Note that our definition implies directly that

$$\mu_0(Lf) = \langle \mathrm{Id}_m^{-1} \Delta f, 1 \rangle = 0, \quad \forall f \in H^1\left((0,1)^N\right),$$

and that $\operatorname{Id}_m^{-1} \Delta f = \Delta f$ in $\mathcal{D}'((0,1)^N)$.

Theorem 3.6. The space V and hence \tilde{V} are densely and compactly embedded in H.

Proof. Denote by \tilde{H} the closure of V in H. To prove that

$$H = H,$$

it then suffices to show that any $f \in H$ orthogonal to V is zero. Let $f \in H$ be orthogonal to V, i.e.,

$$(f|g)_{H^{-1}(T^N)} = 0 \ \forall g \in V.$$

As $\mu_0(f) = \mu_0(g) = 0$, we deduce that

$$\int_{(0,1)^N} \nabla u_f \cdot \nabla \bar{u}_g \, dx = 0 \,\,\forall g \in V.$$

According to (3.1), we get equivalently

$$\langle f, u_q \rangle = 0 \ \forall g \in V.$$

But we will show below that

(3.5)
$$\{u_g : g \in V\} + \operatorname{Har}(T^N) \text{ is dense in } H^1(T^N),$$

and therefore as $f \in H$, we deduce that f = 0 because

$$\langle f, v \rangle = 0 \ \forall v \in H^1\left(T^N\right).$$

It remains to prove (3.5). For $u \in H^1(T^N)$, we have already noticed in Lemma 2.2 that $u - R(\gamma_0 u) \in H^1_0((0,1)^N)$, therefore there exists a sequence of $\psi_n \in \mathcal{D}((0,1)^N)$ such that

$$\psi_n \to u - R(\gamma_0 u)$$
 in $H_0^1((0,1)^N)$ as $n \to \infty$,

or equivalently

(3.6)
$$\psi_n + R(\gamma_0 u) \to u \text{ in } H^1(T^N) \text{ as } n \to \infty.$$

Now for $\psi \in \mathcal{D}((0,1)^N)$, we notice that $L\psi = \Delta \psi \in V$, i.e., $\Delta \psi$ is orthogonal to the (weakly) harmonic functions: In fact if v is a (weakly) harmonic function, then exploiting the fact that ψ has compact support the Gauß–Green formula yields

$$\int_{(0,1)^N} \Delta \psi \bar{v} dx = - \int_{(0,1)^N} \nabla \psi \nabla \bar{v} dx = 0,$$

where the last identity follows by definition of (weakly) harmonic function. Furthermore, $u_{\Delta\psi} = \psi - \mu_0(\psi)$ by Remark 3.4.

Applying these last remarks to (3.6) means that

$$u_{\Delta\psi_n} + \mu_0(\psi_n) + R(\gamma_0 u) \to u \text{ in } H^1(T^N) \text{ as } n \to \infty.$$

As $\mu_0(\psi_n) + R(\gamma_0 u) \in \text{Har}(T^N)$ and $\Delta \psi_n \in V$, the density assertion in (3.5) follows. Finally, compactness follow by the compactness of the embedding $H^1(T^N) \hookrightarrow L^2((0,1)^N)$.

We are finally in the position to prove a formula that can be seen as an H^{-1} analogue of the usual Gauß–Green-formula that hold with respect to the inner product of L^2 .

Lemma 3.7. For all $f \in H^1(T^N)$ and all $h \in L^2((0,1)^N)$ one has

(3.7)
$$(\mathrm{Id}_m^{-1}(\Delta f)|h)_{H^{-1}(T^N)} = -(f|h)_{L^2(T^N)} + (R\gamma_0 f|h)_{L^2(T^N)}.$$

Proof. For all $f \in H^1(T^N)$, as Δf belongs to $H^{-1}((0,1)^N)$, we can set $g = \mathrm{Id}_m^{-1}(\Delta f)$ that satisfies $\mu_0(g) = 0$ and consequently for all $h \in L^2((0,1)^N)$ one has

(3.8)
$$(g|h)_{H^{-1}(T^N)} = \int_{(0,1)^N} \nabla u_g \cdot P_N h \, dx,$$

with

$$P_N h = \nabla u_h + \frac{\mu_0(h)}{N} \left(\vec{x} - \frac{\vec{1}}{2} \right).$$

Owing to (3.1), we get

$$\int_{(0,1)^N} \nabla u_g \cdot \nabla u_h \, dx = -\langle g, u_h \rangle,$$

and by the definition of Id_m^{-1} , we obtain

$$\int_{(0,1)^N} \nabla u_g \cdot \nabla u_h \, dx = -\langle \Delta f, u_h - R\gamma_0 u_h \rangle = -\langle \Delta (f - R\gamma_0 f), u_h - R\gamma_0 u_h \rangle.$$

Hence by the definition of $\Delta(f - R\gamma_0 f)$ as element of $H^{-1}((0,1)^N)$, we get

$$\int_{(0,1)^N} \nabla u_g \cdot \nabla u_h \, dx = \langle \nabla (f - R\gamma_0 f) \cdot \nabla (u_h - R\gamma_0 u_h) \, dx$$
$$= -\int_{(0,1)^N} (f - R\gamma_0 f) \Delta (u_h - R\gamma_0 u_h) \, dx.$$

As $R\gamma_0 u_h$ is harmonic and $\Delta u_h = h - \mu_0(h)$, we get

(3.9)
$$\int_{(0,1)^N} \nabla u_g \cdot \nabla u_h \, dx = -\int_{(0,1)^N} (f - R\gamma_0 f)(h - \mu_0(h)) \, dx$$
$$= -\int_{(0,1)^N} fh \, dx + \int_{(0,1)^N} R\gamma_0 fh \, dx + \mu_0(h) \int_{(0,1)^N} (f - R\gamma_0 f) \, dx.$$

On the other hand, one sees that

$$\int_{(0,1)^N} \nabla u_g \cdot \left(\vec{x} - \frac{\vec{1}}{2} \right) \, dx = \int_{(0,1)^N} \nabla u_g \cdot \nabla v \, dx,$$

where $v \in H^1(T^N)$ is defined by

$$v(x) := \frac{1}{2} \left\| \vec{x} - \frac{\vec{1}}{2} \right\|_{L^2}^2.$$

Hence (3.2) yields

$$\int_{(0,1)^N} \nabla u_g \cdot \left(\vec{x} - \frac{\vec{1}}{2}\right) \, dx = -\langle g, v \rangle,$$

and again by definition of Id_m^{-1} ,

$$\int_{(0,1)^N} \nabla u_g \cdot \left(\vec{x} - \frac{\vec{1}}{2}\right) \, dx = -\langle \Delta f, v - R\gamma_0 v \rangle = -\langle \Delta (f - R\gamma_0 f), v - R\gamma_0 v \rangle.$$

As before we then obtain

$$\int_{(0,1)^N} \nabla u_g \cdot \left(\vec{x} - \frac{\vec{1}}{2} \right) \, dx = -\langle f - R\gamma_0 f, \Delta(v - R\gamma_0 v) \rangle = -N \langle f - R\gamma_0 f, 1 \rangle.$$

This identity and (3.9) in (3.8) yields the conclusion.

Remark 3.8. Let us comment on the special case N = 1. Then for all $h \in L^2(0, 1)$ P_1h is equal to Ph defined as in [16, (2.5)]. Furthermore, also in view of Remark 2.1 we find that V and \tilde{V} agree with the two spaces with same name introduced in [16, § 2]. Therefore, the theory we develop in the present paper is a generalization of that introduced in [16].

4. Operators in the space of zero mean functions

In this section we want to determine precisely two relevant realizations of the Laplacian in H, the space of those functionals that annihilate periodic and (weakly) harmonic functions.

We are going to consider the sesquilinear form a defined by

(4.1)
$$a(f,g) := \int_{(0,1)^N} f(x)\overline{g}(x) \, dx,$$

with form domain either \tilde{V} or V. Since both \tilde{V} and V are dense in H by Lemma 3.6, the form a with domain \tilde{V} or V is associated with a linear operator $(\tilde{A}, D(\tilde{A}))$ or (A, D(A)), respectively, defined by

$$\begin{split} D(\tilde{A}) &:= \left\{ f \in \tilde{V} : \exists g \in H : a(f,h) = \int_{(0,1)^N} \nabla u_g(x) \cdot \nabla \bar{u}_h(x) \, dx \ \forall h \in \tilde{V} \right\}, \\ \tilde{A}f &:= g \end{split}$$

and

$$D(A) := \left\{ f \in V : \exists g \in H : a(f,h) = \int_{(0,1)^N} \nabla u_g(x) \cdot \nabla \bar{u}_h(x) \, dx \ \forall h \in V \right\},$$

$$Af := g.$$

Let us describe these two operators more precisely.

Theorem 4.1. One has

$$D(\tilde{A}) = \tilde{V} \cap (H_m^1(T^N) + V_0),$$

$$\tilde{A}f = -\mathrm{Id}_m^{-1}\Delta f, \quad \forall f \in D(\tilde{A}).$$

Observe that in view of Remark 2.5, each function in $D(\tilde{A})$ is weakly periodic.

Proof. Denote

$$\mathcal{K} := \tilde{V} \cap (H_m^1(T^N) + V_0).$$

Let us first show the inclusion $D(\tilde{A}) \subset \mathcal{K}$. Let $f \in D(\tilde{A})$. Then there exists $g = Af \in H_m^{-1}(T^N)$ (because $g \in H$) such that

$$\int_{(0,1)^N} f(x)\overline{h}(x) \, dx = \int_{(0,1)^N} \nabla u_g(x) \cdot \nabla \overline{u}_h(x) \, dx, \qquad \forall h \in \tilde{V}.$$

But according to (3.1) we then have equivalently

$$\int_{(0,1)^N} f(x)\overline{h}(x) \, dx = -\overline{\langle h, u_g \rangle} = -\int_{(0,1)^N} \overline{h}(x)u_g(x) \, dx, \qquad \forall h \in \tilde{V},$$

because h belongs to $L^2((0,1)^N)$. This means equivalently that $f+u_g$ is orthogonal (in the L^2 sense) to \tilde{V} , i.e., $f+u_g$ belongs to V_0 . As elements of V_0 are harmonic functions, we deduce that

$$\Delta(f+u_g) = 0 \text{ in } \mathcal{D}'\left((0,1)^N\right),$$

and by (3.3), we find

$$\Delta f = -g \text{ in } \mathcal{D}'\left((0,1)^N\right).$$

This shows that Δf belongs to $H^{-1}((0,1)^N)$ and reminding that g belongs to H, we get further

$$g = -\mathrm{Id}_m^{-1}(\Delta f).$$

Similarly as element of V_0 are weakly periodic and $u_g \in H^1(T^N)$ (hence weakly periodic), we deduce that f is weakly periodic.

Let us now prove the converse inclusion. Let $f \in \mathcal{K}$ then $g = -\mathrm{Id}_m^{-1}(\Delta f)$ belongs to H and by Lemma 3.7, we get for any $h \in \tilde{V}$

$$(g|h)_H = (f|h)_{L^2},$$

as $(R\gamma_0 f|h)_{L^2(T^N)} = 0$. This shows that $f \in D(\tilde{A})$ and concludes the proof. \Box

Theorem 4.2. One has

$$D(A) = V \cap (H^1_m(T^N) + V_1),$$

$$Af = -\mathrm{Id}_m^{-1}\Delta f, \quad \forall f \in D(A)$$

Proof. Denote

$$\mathcal{K} := V \cap (H_m^1(T^N) + V_1).$$

Let us first show the inclusion $D(A_0) \subset \mathcal{K}$. Let $f \in D(A_0)$. Then there exists $g =: A_0 f \in H \equiv H_m^{-1}(T^N)$ such that

$$\int_{(0,1)^N} f(x)\overline{h}(x) \, dx = \int_{(0,1)^N} \nabla u_g(x) \cdot \nabla \overline{u}_h(x) \, dx, \qquad \forall h \in V.$$

But according to (3.1) we then have equivalently

$$\int_{(0,1)^N} f(x)\overline{h}(x) \, dx = -\overline{\langle h, u_g \rangle} = -\int_{(0,1)^N} \overline{h}(x)u_g(x) \, dx, \qquad \forall h \in V,$$

because h belongs to $L^2((0,1)^N)$. This means equivalently that $f+u_g$ is orthogonal (in the L^2 sense) to V, i.e., $f+u_g$ belongs to V_1 . As elements of V_1 are harmonic functions, we deduce that

$$\Delta(f+u_g) = 0 \text{ in } \mathcal{D}'\left((0,1)^N\right),$$

and by (3.3), we find

$$\Delta f = -g \text{ in } \mathcal{D}'\left((0,1)^N\right).$$

This shows that Δf belongs to $H^{-1}((0,1)^N)$ and reminding that g belongs to H, we get further

$$g = -\mathrm{Id}_m^{-1}(\Delta f).$$

This shows the desired inclusion because by definition $D(A_0) \subset V$.

The converse inclusion is proved as in the previous Theorem by using Lemma 3.7. $\hfill \square$

Remark 4.3. If one wants to study the Laplacian acting on functions on a more general domain $\Omega \subset \mathbb{R}^N$, one may look for a Lipschitz-homeomorphism mapping $(0,1)^N$ into Ω . In this way, however, also the Laplacian on $(0,1)^N$ is transformed into a different elliptic operator on Ω . The following is a possibly more convenient approach: Take a "partition" $\Gamma = \{\Gamma_1, \ldots, \Gamma_{2N}\}$ of $\partial\Omega$ such that

- each set $\Gamma_i \subset \partial \Omega$ is open,
- the sets Γ_i are pairwise disjoint,
- the union of their closures covers $\partial \Omega$, and

• such that each Γ_i is bijective (say, via some θ_i) to Γ_{N+i} for all i = 1, ..., N. Define

$$H^{1}_{\Gamma}(\Omega) := \{ u \in H^{1}((\Omega)) : \gamma_{i} u = \gamma_{N+i} u \circ \theta_{i} \text{ on } \Gamma_{i}, \forall i = 1, \dots, N \},\$$

where γ_i is the trace operator on Γ_i . In this way, we can regard in a natural way the space $H^1(T^N)$ as the space $H^1_{\Gamma}((0,1)^N)$ with the partition Γ made of faces of the hypercube, and deduce that all the constructions of this paper can be extended to consider realizations (with linear conditions on the moments of order 0 and 1) of the Laplacians acting on functions over general domains Ω .

10

5. Well-posedness of the parabolic problem in the space of zero Mean functions

In the previous section we have showed that the densely defined sesquilinear, symmetric, form a is bounded and coercive both on \tilde{V} and whenever restricted to V. Hence, from the general theory of forms we deduce that the associated operators \tilde{A} and A generate a cosine operator function on H. Since we are especially interested in diffusion-type problems, we want to mention explicitly the following immediate consequence, where we are using the compact embedding of $H^1((0,1)^N)$ in $L^2((0,1)^N)$.

Proposition 5.1. Both operators -A and $-\tilde{A}$ generate analytic, contractive, exponentially stable semigroups.

Hence, we can say that the abstract Cauchy problems associated with A and A are well-posed. However, the challenge is now to understand just *which* are these abstract Cauchy problem. The characterization of D(A) and $D(\tilde{A})$ in Theorem 4.1 and Theorem 4.2 is not very satisfying – and correspondingly poor is the description of the differential equations effectively solved by the semigroups. Yet, we are able to deduce well-posedness of the classical heat equation in the following weak sense.

Theorem 5.2. Let $u_0 \in D(A)$. Then the function

$$t \mapsto u(t, \cdot) := e^{-tA} u_0(\cdot)$$

solves

$$\frac{\partial u}{\partial t}(t,x) = \Delta u(t,x), \qquad u(0) = u_0,$$

with conditions

(5.1)
$$\int_{(0,1)^N} u(t,y)\bar{h}(y)dy = 0 \quad \forall h \in \operatorname{Har}\left((0,1)^N\right), \qquad t > 0,$$

in the sense of distributions, i.e., for all t > 0 (5.1) is satisfied and moreover

$$\left\langle \frac{\partial u}{\partial t}(t,\cdot) - \Delta u(t,\cdot), g \right\rangle_{H-H'} = 0 \quad \text{for all } g \in H^1_0\left((0,1)^N\right).$$

Proof. It suffices to take into account Theorem 4.2 and recall that the analytic semigroup $(e^{-tA})_{t\geq 0}$ maps u_0 into the domain of D(A) for all t > 0, and moreover u solves

$$\left\langle \frac{\partial u}{\partial t}(t,\cdot) - \mathrm{Id}_m^{-1} \Delta u(t,\cdot), g \right\rangle_{H-H'}$$
 for all $g \in H'$

Now, $\operatorname{Har}(T^N)$ is a closed subspace of $H^1(T^N)$ and H is its annihilator, we can deduce that there is an isomorphism

$$\mathfrak{J}: H^1\left(T^N\right)/\mathrm{Har}(T^N) \to H'$$

given by

$$\langle \varphi, \mathfrak{J}[h] \rangle_{H-H'} := \langle \varphi, h \rangle_{H^{-1}(T^N) - H^1(T^N)}, \forall \varphi \in H.$$

Then we have that

$$\left\langle \frac{\partial u}{\partial t}(t,\cdot) - \mathrm{Id}_m^{-1} \Delta u(t,\cdot), g \right\rangle_{H-H'} = 0 \quad \text{for all } g \in H^1(T^N),$$

and in particular

$$\left\langle \frac{\partial u}{\partial t}(t,\cdot) - \mathrm{Id}_m^{-1} \Delta u(t,\cdot), g \right\rangle_{H-H'} = 0 \quad \text{for all } g \in \mathcal{D}\left((0,1)^N\right).$$

Now, the assertion follows from Remark 3.5.

Similarly, exploiting instead Theorem 4.1, we can obtain the following.

Theorem 5.3. Let $u_0 \in D(\tilde{A})$. Then the function

0

$$t \mapsto u(t, \cdot) := e^{-tA} u_0(\cdot)$$

solves

$$\frac{\partial u}{\partial t}(t,x) = \Delta u(t,x), \qquad t > 0, \qquad u(0) = u_0,$$

in the sense of distributions, it is periodic in the sense of Definition 2.3 and satisfies conditions

(5.2)
$$\int_{(0,1)^N} u(t,y)\overline{h}(y)dy = 0 \quad \forall h \in \operatorname{Har}(T^N), \qquad t > 0.$$

Observe that in particular the constant functions belong to $\operatorname{Har}(T^N)$, and therefore H contains all mean zero functions – i.e., all functions with vanishing moment of order 0. Thus, if $f \in \tilde{V}$ (and in particular if $f \in D(\tilde{A})$), then

$$\int_{(0,1)^N} f(x) dx = 0.$$

Furthermore, the functions $g_i : (0,1)^N \ni x = (x_1, \ldots, x_n) \mapsto x_i \in (0,1)$ belong to Har $((0,1)^N)$ for all $i = 1, \ldots, N$. Hence: If additionally $f \in V$ (and in particular if $f \in D(A)$), then

$$\int_{(0,1)^N} f(x) dx = \int_{(0,1)^N} x_i f(x) dx = 0, \qquad i = 1, \dots, N.$$

In particular, for all t > 0 both $(e^{-tA})_{t\geq 0}$ and $(e^{-t\tilde{A}})_{t\geq 0}$ map any $u_0 \in H$ into a function that has mean zero. Furthermore $(e^{-tA})_{t\geq 0}$ maps any $u_0 \in H$ into a function that has vanishing first linear moments along each axis, as such functions are referred to in [19, § 9.6.5].

6. L^2 -results and interplays with extension theory

It is well-known that if an operator that comes from a sesquilinear form generates an analytic semigroup, then so does its part in the form domain. In this conclusive section we will devote our attention in particular to the part in V of the operator A described in Theorem 4.2. To begin with, we recall a celebrated result due to M. Krein: Each closed, densely defined, symmetric, positive definite operator has one smallest self-adjoint positive definite extension¹. This extension is nowadays commonly called the *Krein–von Neumann extension* – cf. [2] for a brief introduction to this subject including the connection with the so-called "buckling problem" of mathematical physics, or [3] and [18, Chapter 13] for very comprehensive overviews.

It turns out that such an extension can be also characterized as follows – this is but a special instance of [3, Thm. 2.7].

Lemma 6.1. Let S be a closed, densely defined, symmetric operator on a Hilbert space $\mathcal{H} = L^2(\Omega)$ such that $S - \epsilon$ Id is positive definite for some $\epsilon > 0$. Among all its self-adjoint extensions there exists exactly one whose domain contains the null space of S^* : This is precisely the Krein–von Neumann extension.

Now, let S be the Laplacian on $L^2(T^N)$ defined on test functions only,

$$Sf := -\Delta f, \qquad f \in \mathcal{D}((0,1)^N)$$

Then, S is clearly symmetric (but of course not self-adjoint) and negative definite, and in fact it follows from the Poincaré inequality that $S - \epsilon$ Id is positive definite for some $\epsilon > 0$ (for example, if one takes ϵ to be the lowest eigenvalue of the

¹ In the sense of forms: Let a_1, a_2 be two densely defined symmetric bounded elliptic forms with associated operators A_1, A_2 . If their form domains $D(a_1), D(a_2)$ satisfy $D(a_2) \subset D(a_1)$ and $a_1(x, x) \leq a_2(x, x)$ for all $x \in D(a_2)$, then A_1 is said to be smaller than A_2 .

Laplacian with Dirichlet boundary conditions). It follows from our construction that the domain of the part A_V of A in the form domain V, i.e.,

$$D(A_V) = \{ v \in V : Av \in V \}$$

is orthogonal to the null space of S^\ast (that is, to the space of (weakly) harmonic functions.

Let A_K be the extension of A_V defined by

$$D(A_K) = D(A_V) \oplus V_1,$$

and

$$A_K v = A_V v_0, \quad \forall v = v_0 \oplus v_1 \in D(A_V) \oplus V_1.$$

Hence A_K is a selfadjoint operator in $L^2((0,1)^N)$, and therefore $-A_K$ is exactly the Krein–von Neumann extension of S.

We thus obtain the following alternative characterization of the Laplacian under conditions on the moments of order 0 and 1.

Theorem 6.2. The L^2 -realization $-A_V$ of the Laplacian agrees with the reduced Krein–von Neumann Laplacian as introduced in [3, § 2]. In particular,

$$A_V f = \Delta f, \qquad \forall f \in D(A_V),$$

(and not only $A_V f = \operatorname{Id}_m^{-1} \Delta f!$).

Furthermore, because $\Omega = T^N$ is a bounded Lipschitz domain (hence a quasiconvex domain in the sense of [3, § 5.1]) one can apply the general theory surveyed in [3] and in particular by [3, Thm 6.5] $-A_K$ is precisely the Laplacian with boundary conditions given by

$$\frac{\partial u}{\partial \nu} = \mathbf{D} \mathbf{N} u_{|\partial \Omega} \qquad \text{on } \partial \Omega,$$

where **D**N is the Dirichlet-to-Neumann operator associated with Δ on $\partial\Omega$, defined in a suitably weak sense, see [3, § 5–6].

Remark 6.3. In case of N = 1 we recover in particular the boundary conditions

$$u'(1) = u'(0) = u(1) - u(0)$$

This in accordance with [16, Cor. 3.10] and also with the setting of [4]. The same boundary conditions are also referred to as *transparent conditions* in numerical analysis and δ' -interaction in mathematical physics, cf. respectively [12, 9] for an interpretation of these conditions and a list of related references. While the Krein–von Neumann extension is already known to have interesting connections with elasticity theory, cf. [3], the observation that the associated heat equation enjoys conservation of moments of order 0 and 1 seems to be new in the literature.

References

- W. Arendt and T. ter Elst. Sectorial forms and degenerate differential operators. J. Operator Th., 67:33-72, 2012.
- [2] M. Ashbaugh, F. Gesztesy, M. Mitrea, R. Shterenberg, and G. Teschl. The Krein–von Neumann extension and its connection to an abstract buckling problem. *Math. Nachr.*, 283:165– 179, 2010.
- [3] M. Ashbaugh, F. Gesztesy, M. Mitrea, R. Shterenberg, and G. Teschl. A Survey on the Kreinvon Neumann Extension, the Corresponding Abstract Buckling Problem, and Weyl-type Spectral Asymptotics for Perturbed Krein Laplacians in Nonsmooth Domains. In M. Demuth and W. Kirsch, editors, *Mathematical Physics, Spectral Theory and Stochastic Analysis*, pages 1–106, Berlin, 2013. Springer-Verlag.
- [4] A. Bobrowski and D. Mugnolo. On moment preserving cosine families and semigroups in C[0,1]. J. Evol. Equ., (in press):DOI 10.1007/s00028-013-0199-x, 2013.
- [5] A. Bouziani and N. Benouar. Problème mixte avec conditions intégrales pour une classe d'équations hyperboliques. Bull. Belg. Math. Soc., 3:137–145, 1996.
- [6] J. Cannon. The solution of the heat equation subject to the specification of energy. Quart. Appl. Math, 21:155–160, 1963.

DELIO MUGNOLO AND SERGE NICAISE

- [7] J. Cannon. The One-Dimensional Heat Equation, volume 23 of Encyclopedia of Mathematics and its Applications. Cambridge Univ. Press, Cambridge, 1984.
- [8] F. Demengel and G. Demengel. Functional Spaces for the Theory of Elliptic Partial Differential Equations. Universitext. Springer-Verlag, Berlin, 2007.
- [9] P. Exner. Contact interactions on graph superlattices. J. Phys. A, 29:87–102, 1996.
- [10] G. Greiner. Perturbing the boundary conditions of a generator. Houston J. Math, 13:213–229, 1987.
- [11] P. Grisvard. Elliptic Problems in Nonsmooth Domains, volume 24 of Monographs and Studies in Mathematics. Pitman, Boston, MA, 1985.
- [12] L. Halpern and J. Rauch. Absorbing boundary conditions for diffusion equations. Numer. Math., 71:185–224, 1995.
- [13] J. Lions and E. Magenes. Non-Homogeneous Boundary Value Problems and Applications, volume 181–183 of Grundlehren der mathematischen Wissenschaften. Springer-Verlag, Berlin, 1972.
- [14] N. Merazga and A. Bouziani. Rothe method for a mixed problem with an integral condition for the two-dimensional diffusion equation. *Abstr. Appl. Anal.*, 16:899–922, 2003.
- [15] S. Mesloub and A. Bouziani. Mixed problem with integral conditions for a certain class of hyperbolic equations. J. Appl. Math., 1:107–116, 2001.
- [16] D. Mugnolo and S. Nicaise. Well-posedness and spectral properties of heat and wave equations with non-local conditions. J. Differ. Equ., (in press), 2013.
- [17] L. Pul'kina. Nonlocal problem with a first-kind integral condition for a multidimensional hyperbolic equation. Dokl. Math., 76:741–743, 2007.
- [18] K. Schmüdgen. Unbounded Self-adjoint Operators on Hilbert Space, volume 265 of Graduate Texts in Mathematics. Springer-Verlag, Berlin, 2012.
- [19] J. Vázquez. Smoothing and Decay Estimates for Nonlinear Diffusion Equations Equations of Porous Medium Type, volume 33 of Oxford Lect. Ser. Math. and its Appl. Oxford Univ. Press, Oxford, 2006.

DELIO MUGNOLO, INSTITUT FÜR ANALYSIS, UNIVERSITÄT ULM, 89069 ULM, GERMANY *E-mail address*: delio.mugnolo@uni-ulm.de

SERGE NICAISE, UNIVERSITÉ DE VALENCIENNES ET DU HAINAUT CAMBRÉSIS, LAMAV, FR CNRS 2956, ISTV, F-59313 - VALENCIENNES CEDEX 9, FRANCE

E-mail address: Serge.Nicaise@univ-valenciennes.fr